

## **BDNF Val66Met moderates memory impairment, hippocampal function and tau in preclinical autosomal dominant Alzheimer's disease**

Yen Ying Lim<sup>1</sup>, Jason Hassenstab<sup>2</sup>, Carlos Cruchaga<sup>3</sup>, Alison Goate<sup>4</sup>, Anne M. Fagan<sup>2</sup>, Tammie L.S. Benzinger<sup>5</sup>, Paul Maruff<sup>1,6</sup>, Peter J. Snyder<sup>7</sup>, Colin L. Masters<sup>1</sup>, Ricardo Allegri<sup>8</sup>, Jasmeer Chhatwal<sup>9,10</sup>, Martin R. Farlow<sup>11</sup>, Neill R. Graff-Radford<sup>12</sup>, Christoph Laske<sup>13,14</sup>, Johannes Levin<sup>15</sup>, Eric McDade<sup>2</sup>, John M. Ringman<sup>16</sup>, Martin Rossor<sup>17</sup>, Stephen Salloway<sup>7</sup>, Peter R. Schofield<sup>18,19</sup>, David M. Holtzman<sup>2</sup>, John C. Morris<sup>2</sup>, and Randall J. Bateman<sup>2</sup>, on behalf of the Dominantly Inherited Alzheimer Network

1. The Florey Institute, The University of Melbourne, Parkville, Victoria, Australia
2. Department of Neurology, Washington University in St. Louis, St. Louis, MO, USA
3. Department of Psychiatry, Washington University in St. Louis, St. Louis, MO, USA
4. Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, USA
5. Department of Radiology, Washington University in St. Louis, St. Louis, MO, USA
6. Cogstate Ltd., Melbourne, Victoria, Australia
7. Department of Neurology, Warren Alpert Medical School of Brown University, Providence, RI, USA
8. Ageing and Memory Center, Instituto de Investigaciones Neurologicas "Raúl Carrea" (FLENI), Buenos Aires, Argentina
9. Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
10. Center for Alzheimer Research and Treatment, Department of Neurology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA
11. Department of Neurology, Indiana University School of Medicine, Indianapolis, IN, USA
12. Department of Neurology, Mayo Clinic Jacksonville, FL, USA
13. German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany
14. Section for Dementia Research, Department of Cellular Neurology, Hertie Institute for Clinical Brain Research and Department of Psychiatry and Psychotherapy, University of Tübingen, Tübingen, Germany
15. Department of Neurology, University of Munich, Munich, Germany
16. Memory and Aging Center, Keck School of Medicine of the University of Southern California, CA, USA
17. Dementia Research Centre, Department of Neurodegenerative Diseases, UCL Institute of Neurology, London, UK
18. Neuroscience Research Australia, Sydney, NSW, Australia
19. School of Medical Sciences, University of New South Wales, Sydney, NSW, Australia

Address for correspondence:

Yen Ying Lim  
155 Oak Street,  
Parkville, VIC 3052, Australia  
Email: [yen.lim@florey.edu.au](mailto:yen.lim@florey.edu.au)  
Ph: +61 3 9389 2909  
Fax: +61 3 9387 5061

Word Count: 4489 words

---

This is the author's manuscript of the article published in final edited form as:

Lim, Y. Y., Hassenstab, J., Cruchaga, C., Goate, A., Fagan, A. M., Benzinger, T. L. S., ... Bateman, R. J. (2016). BDNF Val66Met moderates memory impairment, hippocampal function and tau in preclinical autosomal dominant Alzheimer's disease. *Brain*, 139(10), 2766–2777. <https://doi.org/10.1093/brain/aww200>

## Abstract

**Background:** The brain-derived neurotrophic factor (*BDNF*) Val66Met polymorphism is implicated in synaptic excitation and neuronal integrity, and has previously been shown to moderate A $\beta$ -related memory decline and hippocampal atrophy in preclinical sporadic Alzheimer's disease (AD). However, the effect of *BDNF* in autosomal dominant AD (ADAD) is unknown. We aimed to determine the effect of *BDNF* Val66Met on cognitive function, hippocampal function, tau and A $\beta$  in preclinical ADAD. We explored effects of apolipoprotein E (*APOE*)  $\epsilon$ 4 on these relationships.

**Methods:** The Dominantly Inherited Alzheimer Network (DIAN) conducted clinical, neuropsychological, genetic, biomarker and neuroimaging measures at baseline in 131 mutation non-carriers (NC) and 143 preclinical ADAD mutation carriers (MC) on average 12 years prior to clinical symptom onset. *BDNF* genotype data were obtained for MCs (95 Val<sub>66</sub> homozygotes, 48 Met<sub>66</sub> carriers).

**Findings:** Among preclinical MCs, Met<sub>66</sub> carriers had worse memory performance, lower hippocampal glucose metabolism and increased levels of CSF tau and phosphorylated tau (p-tau) than Val<sub>66</sub> homozygotes. Cortical A $\beta$  and CSF A $\beta$ <sub>42</sub> levels were significantly different from NC's but did not differ between preclinical MC Val<sub>66</sub> homozygotes and Met<sub>66</sub> carriers. There was an effect of *APOE* on A $\beta$  levels but not cognitive function, glucose metabolism or tau.

**Interpretation:** As in sporadic AD, the deleterious effects of A $\beta$  on memory, hippocampal function, and tau in preclinical ADAD mutation carriers are greater in Met<sub>66</sub> carriers. To date, this is the only genetic factor found to moderate downstream effects of A $\beta$  in ADAD.

## Introduction

Alzheimer's disease (AD) begins with the aggregation of beta-amyloid (A $\beta$ ), the development and spread of hyperphosphorylated tau (Ballatore *et al.* , 2007, Ittner and Götz, 2011), and ultimately neuronal and synaptic loss. This characteristic pathological process manifests initially as cognitive impairment which increases progressively so eventually classification of dementia is warranted (Hardy and Higgins, 1992, Ittner and Götz, 2011, Spiess-Jones and Hyman, 2014). Clinical pathological relationships in AD are still not understood completely, however recent *in vitro* (Hariri *et al.* , 2003, Lee *et al.* , 2012), *post-mortem* (Buchman *et al.* , 2016, Garzon and Fahnestock, 2007, Peng *et al.* , 2005) and animal (Caccamo *et al.* , 2010, Lee *et al.* , 2012, Rosa and Fahnestock, 2015) studies suggest neurotrophic factors, such as brain-derived neurotrophic factor (BDNF) moderate neuronal and synaptic dysfunction and their behavioral expression in AD (Fahnestock, 2011, Lu *et al.* , 2013).

Clinical studies of the role of BDNF in AD are limited by the absence of validated biomarkers for central nervous system (CNS) BDNF (Forlenza *et al.* , 2010, Kim *et al.* , 2015). However, the *BDNF* Val66Met (rs6265) polymorphism Met protein can result in reduced dendritic trafficking and synaptic localization of the protein and up to a 30% reduction in activity-dependent BDNF secretion (Chen *et al.* , 2006, Egan *et al.* , 2003). In healthy young adults, memory dependent hippocampal activity is reduced in Met<sub>66</sub> carriers (Hariri *et al.* , 2003). In the preclinical and prodromal stages of sporadic AD, prospective studies show Met<sub>66</sub> carriers to have increased rates of decline in episodic memory and hippocampal atrophy relative to Val<sub>66</sub> homozygotes (Feng *et al.* , 2013, Lim *et al.* , 2014, Lim *et al.* , 2013, Lim *et al.* , 2014). These same studies observe rates of cortical A $\beta$  accumulation to be unaffected by the Met<sub>66</sub> allele (Lim *et al.* , 2014, Lim *et al.* , 2013), suggesting that *BDNF* Met<sub>66</sub> may accelerate neuronal dysfunction and memory decline by moderating pathological processes downstream of cortical A $\beta$  accumulation, such as tau aggregation.

While the processes that give rise to cortical A $\beta$  accumulation are likely to differ between sporadic and autosomal dominant AD, the effects of A $\beta$  on neurodegeneration and cognition are similar, albeit occurring at markedly younger ages in ADAD (mean age of onset is 45 years) (Bateman *et al.* , 2012, Jack and Holtzman, 2013, Ryman *et al.* ,

2014). Therefore the aim of this study was to investigate the effects of the *BDNF* Met<sub>66</sub> allele on episodic memory, hippocampal function, A $\beta$  and tau in ADAD. The first hypothesis was that in preclinical ADAD mutation carriers (MCs), impairment in episodic memory and hippocampal function would be greater in individuals who carry at least one copy of the *BDNF* Met<sub>66</sub> allele compared to Val<sub>66</sub> homozygotes. The second hypothesis was that cortical A $\beta$  levels would be unrelated to variation in *BDNF* Val66Met. The third hypothesis was that CSF tau levels would be greater in *BDNF* Met<sub>66</sub> carriers compared to Val<sub>66</sub> homozygotes. We also explored the extent to which carriage of the *BDNF* Met<sub>66</sub> allele was associated with domains of cognition beyond episodic memory, neuronal function in the precuneus and CSF biomarkers of A $\beta$ <sub>1-42</sub> and phosphorylated tau (p-tau<sub>181</sub>). Finally, while the apolipoprotein E (*APOE*)  $\epsilon$ 4 allele does not increase severity of clinical presentation in ADAD (Ryman *et al.* , 2014), we observed previously additive effects of the *BDNF* Met<sub>66</sub> and *APOE*  $\epsilon$ 4 alleles on A $\beta$  related cognitive decline in preclinical sporadic AD (Lim *et al.* , 2014). Therefore, we also explored the extent to which *APOE* acts independently, or with *BDNF*, to impact disease processes in ADAD.

## Method

### Participants

Individuals at risk for carrying a mutation for ADAD (i.e., *presenilin 1* [*PSEN1*], *presenilin 2* [*PSEN2*], or *amyloid precursor protein* [*APP*] mutations) were enrolled in the Dominantly Inherited Alzheimer Network (DIAN) study. Participants from families with known pathogenic ADAD mutations were recruited from 197 families at six sites in the United States, one in the United Kingdom and three in Australia (Morris *et al.* , 2012). The process of recruitment and enrollment has been described in detail previously (Bateman *et al.* , 2012, Morris *et al.* , 2012). Baseline data from 274 participants (131 NCs, 143 preclinical MCs) who were cognitively normal, as defined by a Clinical Dementia Rating of 0, and who had completed assessments of cognitive function, neuroimaging and cerebrospinal fluid (CSF) sampling were included. *APOE* genotype was determined for all individuals as part of the DIAN study protocol. Additionally, for MCs, only individuals whose *BDNF* Val66Met polymorphism was available were included. Table 1 shows the demographic characteristics of each participant group.

## **Clinical Assessment**

Without reference to participants' performance on the neuropsychological test battery, a clinician assessed each participant for the presence and severity of clinical symptoms of dementia at baseline. This was operationalized using the Clinical Dementia Rating (CDR) scale, for which a CDR total score of 0 indicates cognitive normality (Morris, 1983). Participants also completed the Mini Mental State Examination (MMSE) and the Geriatric Depression Scale (GDS) at baseline.

## **Neuropsychological Assessment**

All participants were assessed using the DIAN neuropsychological test battery, which includes the Wechsler Memory Scale–Revised Logical Memory (Story A only, immediate and delayed recall) and Digit Span; Category Fluency (animals, vegetables); Trail Making Test A and B; Digit Symbol from the Wechsler Adult Intelligence Scale–Revised (WAIS–R); the Boston Naming Test (30 odd items), letter fluency for F, A, and S, and immediate and delayed recall of a single presentation of a 16-item word list (Storandt *et al.*, 2014). These tasks have been described previously, and were administered according to standard protocols by trained research assistants (Storandt *et al.*, 2014). The process of standardization and quality control of neuropsychological assessments across all DIAN sites have also been described previously (Storandt *et al.*, 2014).

Outcome measures for each neuropsychological test were standardized against the baseline mean and standard deviation for the NC group. Standardized scores were then averaged to form four cognitive domain-specific composite scores for *episodic memory* (Logical Memory delayed recall, word list learning delayed recall); *executive function* (Letter Fluency, Trail Making Test B); *language* (Category Fluency animals + vegetables, Boston Naming Test); *attention* (Digit Span Forwards, Digit Symbol); and *global cognition* (Logical Memory delayed recall, word list learning delayed recall, Digit Symbol, MMSE) (Donohue *et al.*, 2014).

## Genotyping

Genotyping for pathogenic mutations in the *APP*, *PSEN1*, and *PSEN2* genes were performed on DNA extracted from peripheral blood samples using methods described previously (Talbot *et al.*, 1994). Samples were also genotyped with the Infinium HumanExomeCore V1.0 Beadchip (Illumina, Inc). Genotyping was performed at The Genome Technology Access Center (GTAC; <https://gtac.wustl.edu/>) at Washington University. All samples and genotypes underwent stringent quality control (QC). Genotype data were cleaned by applying a minimum call rate for SNPs and individuals (98%). SNPs not in Hardy-Weinberg equilibrium ( $P < 1 \times 10^{-6}$ ) were excluded. No SNPs were removed because low MAF. Gender identification was verified by analysis of X-chromosome SNPs. We tested for unanticipated duplicates using pairwise genome-wide estimates of proportion identity-by-descent using PLINK v1.9. Genotype data for the *BDNF* Val66Met (rs6265) polymorphism was extracted from using PLINK. Clinicians were blinded to all genetic information and genetic polymorphisms were not used diagnostically. *BDNF* Val66Met genotyping was performed only in samples from individuals with a known ADAD mutation.

## Neuroimaging

Images obtained through positron emission tomography (PET) with the use of fluorodeoxyglucose (FDG) and Pittsburgh compound B (PiB) (FDG-PET and PiB-PET, respectively) were co-registered with individual MRI images for region-of-interest (ROI) determination. 3 Tesla volumetric T1-weighted MRI scans from DIAN participants were acquired and processed through FreeSurfer (Martinos Center, Boston, MA) as previously described (Benzinger *et al.*, 2013). Amyloid imaging was performed with a bolus injection of approximately 15 mCi of [11C] PiB. Dynamic imaging acquisition started either at injection for 70 minutes or 40 minutes post-injection for 30 minutes. For analysis, PiB-PET data between 40 to 70 minutes were used. For PiB-PET, total neocortical SUVR was used to determine levels of cortical A $\beta$  deposition, using cerebellar grey matter as the reference region and applying partial volume correction using a regional point spread function as previously described (Su *et al.*, 2015).

Metabolic imaging with [18F] FDG-PET was performed with a 3D dynamic acquisition began 40 minutes after a bolus injection of approximately 5 mCi of FDG and

lasted for 20 minutes. In accordance with previous reports (Bateman *et al.*, 2012), the ROIs selected for this study were the hippocampus and the precuneus, with decreased FDG SUVR indicating decreased glucose metabolism and therefore reduced neuronal function in that area. The reference region used was the cerebellar cortex.

### **Biochemical Analysis**

Fasted CSF was collected in the morning via lumbar puncture. Samples were shipped on dry ice to the DIAN biomarker core laboratory. CSF concentrations of A $\beta$ <sub>42</sub>, total tau, and tau phosphorylated at threonine 181 (ptau<sub>181</sub>) were measured by immunoassay (INNOTEST  $\beta$ -Amyloid1-42, Innogenetics). All values had to meet quality-control standards, including a coefficient of variation of 25% or less, kit “controls” within the expected range as defined by the manufacturer, and measurement consistency between plates of a common sample that was included in each run.

### **Estimated year of onset**

The estimated year from expected symptom onset (EYO) was calculated as the age of the participant at the time of the baseline assessment minus the mean age at onset of all other individuals with the same mutation type (Ryman *et al.*, 2014).

### **Data Analysis**

The study hypotheses that in ADAD, MC *BDNF* Met<sub>66</sub> carriage would be associated with greater impairment in memory and hippocampal function, higher CSF tau but not cortical A $\beta$  levels were tested by submitting the episodic memory composite, PiB-PET A $\beta$ , CSF tau and glucose metabolism in the hippocampus (FDG-PET) to separate analyses of covariance (ANCOVA). In each ANCOVA, EYO was added as a covariate, and Group (NC, MC Val<sub>66</sub>/Val<sub>66</sub>, MC Met<sub>66</sub>) as a fixed factor. Within each ANCOVA, two planned comparisons were constructed with the first comparing MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers and the second comparing MC Val<sub>66</sub> homozygotes to the NC group. Exploratory analyses were conducted only if a statistically significant difference between the MC Val<sub>66</sub> homozygote and MC Met<sub>66</sub> carrier groups was observed for at least one of the primary outcome measures. With this criterion met, the ANCOVAs were repeated for the remaining cognitive composite scores, CSF A $\beta$ <sub>42</sub>, CSF p-

tau<sub>181</sub>, and FDG-PET in the precuneus. The extent to which the presence of the *APOE*  $\epsilon$ 4 allele influenced the effect of *BDNF* on cognitive function, A $\beta$  burden, tau and neuronal function was determined by repeating these analyses with  $\epsilon$ 4 status (carrier vs. non-carriers) entered into all statistical models. Finally, to further understand the effect of *BDNF* Val66Met on cognitive and biomarker outcomes in ADAD, we expressed each cognitive and biomarker outcome variable as a function of EYO. For the primary outcomes, statistical significance was classified as  $p < .05$ . This was to balance the risk of false positive findings against the identification of important relationships because (a) this is an exploratory investigation in a relatively new area in which an important clinical issue has been identified, (b) as all four primary outcome measures are recognized as part of the AD pathological process, changes in these will be correlated and (c) effect sizes (Cohen's  $d$ ) were used to guide interpretation about the meaningfulness of statistical tests and comparisons with effect sizes  $< 0.2$  were classified as trivial and not interpreted regardless of statistical significance (Cohen, 1988).

## Results

### Demographic and clinical characteristics

MCs were significantly younger than NCs, although the EYO between MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers did not differ significantly. NC and MC groups did not differ on any other demographic characteristic. While the inclusion criteria required all individuals to have a CDR score of 0, the CDR sum of boxes score was significantly higher in MC Met<sub>66</sub> carriers than in MC Val<sub>66</sub> homozygotes and NCs (Table 1). Groups did not differ in MMSE total scores or levels of depressive symptoms.

### Effect of *BDNF* Val66Met on episodic memory, cortical A $\beta$ , CSF tau and glucose metabolism in the hippocampus

Group means and standard deviations for raw scores on each of the primary outcome cognitive and biomarker measures for each group are summarized on Table 2. The outcomes of the primary analyses are summarized on Figure 1 for episodic memory and Figure 2 for the AD biomarkers. Statistically significant group differences between MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers were observed for episodic memory (Figure 1), glucose metabolism in the hippocampus and CSF tau, but not cortical A $\beta$



(Figure 2). Effect sizes for these comparisons were, by convention, moderate-to-large in magnitude for episodic memory, glucose metabolism in the hippocampus and CSF tau levels, but were trivial for levels of cortical A $\beta$ . No statistically significant differences between NC and MC Val<sub>66</sub> homozygotes were observed for any of the primary outcome measures, with all differences small in magnitude.

### **Effect of *BDNF* Val66Met on cognition, CSF A $\beta$ <sub>42</sub>, CSF p-tau<sub>181</sub> and glucose metabolism in the precuneus**

For each exploratory cognitive and biomarker outcome measure, raw group means and standard deviations are summarized on Table 2. Figure 1 and 2 also summarises the outcomes of the exploratory analyses for cognitive measures and AD biomarkers respectively. Statistically significant group differences, of a moderate-to-large magnitude, were observed between MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers for CSF p-tau<sub>181</sub> levels (Figure 2, exploratory outcomes) but not for glucose metabolism in the precuneus or for the executive function, language, attention or global cognition composites (Figure 1, exploratory outcomes). There were also no statistically significant differences between MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers on CSF A $\beta$ <sub>42</sub> levels, with these differences small in magnitude (Figure 2, exploratory outcomes).

When compared to NCs, MC Val<sub>66</sub> homozygotes showed no statistically significant impairment in any domain of cognitive function (Figure 1) and did not differ significantly in the extent of glucose metabolism in the hippocampus or the precuneus (Figure 2). Compared to NCs, both MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers showed elevated levels of CSF tau and p-tau<sub>181</sub>, and increased PiB-PET SUVR and decreased CSF A $\beta$ <sub>42</sub> levels (Table 2).

### **Effect of *APOE* $\epsilon$ 4 on cognitive function, neuronal dysfunction, A $\beta$ and tau**

Re-analyses of the primary hypotheses with the addition of *APOE* status indicated no significant main effect of *APOE* status and no significant interaction between *APOE* and *BDNF* status on any measure of cognitive function (Table 3). Similarly, there was no significant main effect of *APOE* or interaction between *APOE* and *BDNF* for any outcome measure of glucose metabolism or tau (Table 3). However, there was a significant main effect of *APOE* for both PiB-PET SUVR and CSF A $\beta$ <sub>42</sub>, although

there were no significant interactions between *APOE* and *BDNF* for either measure (Table 3). Post-hoc analyses showed that when compared to MC  $\epsilon 4$  non-carriers, MC  $\epsilon 4$  carriers had significantly increased PiB-PET SUVR ( $d$  [95%CI = 0.45 [0.05, 0.85],  $p = .03$ ) and decreased CSF  $A\beta_{42}$  levels ( $d$  [95%CI = 0.76 [0.34, 1.17],  $p < .001$ ).

### **Effect of *BDNF* Val66Met on the relationship between EYO and markers of cognitive and neuronal function, $A\beta$ and tau**

There were no statistically significant relationships between level of cognitive function and EYO in NCs or in MC Val<sub>66</sub> homozygotes. However, the relationship between EYO and episodic memory was statistically significant and moderate in magnitude for MC Met<sub>66</sub> carriers (Figure 3A). Similarly, there were no statistically significant relationships between glucose metabolism in the hippocampus and EYO in NCs or in MC Val<sub>66</sub> homozygotes. However, the relationship between glucose metabolism in the hippocampus and EYO was moderate in magnitude and statistically significant for MC Met<sub>66</sub> carriers (Figure 3B).

There was no relationship between levels of cortical  $A\beta$  and EYO in NCs, but there was a significant moderate association between cortical  $A\beta$  levels and EYO in MCs irrespective of *BDNF* Val66Met polymorphism (Figure 3C). Similarly, while there was no association between CSF tau and EYO in NCs, there was a significant moderate association between CSF tau and EYO in MCs, irrespective of *BDNF* Val66Met genotype (Figure 3D), with MC Met<sub>66</sub> carriers showing systematically higher levels of CSF tau relative to their EYO than MC Val<sub>66</sub> homozygotes (Figure 3D).

## **Discussion**

The results show that the presence of one copy of the *BDNF* Met<sub>66</sub> allele increased the severity of impairment in episodic memory and hippocampal function in preclinical ADAD. This effect is clinically important as the magnitude of memory impairment related to MC Met<sub>66</sub> carriers was approximately double that observed in MC Val<sub>66</sub> homozygotes. These findings in the DIAN cohort are consistent with the greater memory decline and hippocampal volume loss observed in older adults with preclinical or prodromal sporadic AD from the AIBL and ADNI studies (Feng *et al.*, 2013, Lim *et al.*,

2014, Lim *et al.* , 2013). The results confirm therefore, in an independent sample, that *BDNF* is important to the preclinical presentation of AD.

The current data support the first hypothesis that in preclinical MCs, impairment in memory and hippocampal function would be greater in Met<sub>66</sub> carriers compared to Val<sub>66</sub> homozygotes. Compared to MC Val<sub>66</sub> homozygotes, MC Met<sub>66</sub> carriers had worse episodic memory function (Figure 1). In contrast, no memory impairment was observed in MC Val<sub>66</sub> homozygotes compared to NCs. Similarly, hippocampal function, determined by cerebral glucose metabolism, was also reduced in MC Met<sub>66</sub> carriers compared to MC Val<sub>66</sub> homozygotes. However, MC Val<sub>66</sub> homozygotes did not show lower glucose metabolism compared to NCs. As increased oxidative stress has been previously observed in females (Keaney *et al.* , 2003), it is possible that the sex of participants may better account for the memory impairment in MC Met<sub>66</sub> carriers. However, re-analysis of all primary outcome measures suggest that even when the sex of participants was considered, the effect of *BDNF* Val66Met on memory impairment, hippocampal function and tau remains (Table 4). Finally, MC Met<sub>66</sub> carriers who were estimated to be nearer to their expected year of clinical symptom onset (EYO) showed increased memory impairment and lower glucose metabolism in the hippocampus (Figure 3). In contrast, EYO was not associated with memory impairment or glucose metabolism in NCs or MC Val<sub>66</sub> homozygotes.

The second hypothesis that cortical A $\beta$  and CSF A $\beta$ <sub>42</sub> levels would be unrelated to allelic variation in *BDNF* Val66Met was also supported. Preclinical MC Met<sub>66</sub> carriers and Val<sub>66</sub> homozygotes had equivalent levels of higher cortical A $\beta$  and CSF A $\beta$ <sub>42</sub>. Furthermore, these group differences were, by convention, small (i.e.,  $d < 0.2$ ; Figure 2) in magnitude indicating that absence of statistically significant differences was not due to insufficient statistical power. Compared to NCs, both Met<sub>66</sub> carriers and Val<sub>66</sub> homozygotes showed increased levels of cortical A $\beta$  deposition and decreased levels of CSF A $\beta$ <sub>42</sub>. Similarly, cortical A $\beta$  burden was higher in preclinical MCs who were nearer to their EYO; although this relationship was not moderated by the *BDNF* Val66Met polymorphism (Figure 3C). Increased cortical A $\beta$  and lower CSF A $\beta$ <sub>42</sub> levels have been observed previously in preclinical ADAD (Bateman *et al.* , 2012, Ryman *et al.* , 2014). The absence of any effect of Met<sub>66</sub> carriage on A $\beta$  burden in preclinical ADAD is also consistent with observations that *Met* carriage was unrelated to rates of cortical A $\beta$

accumulation over 3 years in preclinical and prodromal sporadic AD (Feng *et al.* , 2013, Lim *et al.* , 2014, Lim *et al.* , 2013). Together, these findings suggest that the effect of the *BDNF* Met<sub>66</sub> allele is independent of the effect of A $\beta$  on risk for, and progression of, AD.

The results also support the third hypothesis that CSF levels of tau would be greater in MC Met<sub>66</sub> carriers compared to MC Val<sub>66</sub> homozygotes. Levels of both CSF tau and p-tau<sub>181</sub> were increased substantially in preclinical MC Met<sub>66</sub> carriers compared to preclinical MC Val<sub>66</sub> homozygotes (Figure 2). Compared to NCs, preclinical MC Val<sub>66</sub> homozygotes also showed increased levels of CSF tau and p-tau<sub>181</sub>, although not to the same extent as MC Met<sub>66</sub> carriers. Despite the overall increase in these biochemical markers, strong relationships between EYO and CSF tau were observed in both MC Val<sub>66</sub> homozygotes and MC Met<sub>66</sub> carriers, and the magnitude of these relationships were equivalent (Figure 3D). Thus, while the Met<sub>66</sub> allele hastens memory dysfunction in preclinical ADAD, it does not necessarily affect the rate at which p-tau<sub>181</sub> accumulates in CSF. Instead, substantial differences in CSF p-tau<sub>181</sub> levels between MC Met<sub>66</sub> carriers and MC Val<sub>66</sub> homozygotes (Figure 2) suggest that MC Val<sub>66</sub> homozygotes may have an increased level of resilience to the neurotoxic effects of tau and A $\beta$ .

Finally, we explored the extent to which *APOE* acts independently or with *BDNF* to impact disease processes in ADAD. There were no independent effects of *APOE*  $\epsilon$ 4, or combined effects of *APOE* and *BDNF*, on cognition, neuronal function or CSF tau (Table 3). However, compared to MC  $\epsilon$ 4 non-carriers, MC  $\epsilon$ 4 carriers showed increased cortical A $\beta$  and decreased CSF A $\beta$ <sub>42</sub>. This indicates that in preclinical ADAD, the abnormal accumulation of cortical A $\beta$  resulting from pathogenic mutations is increased further by the *APOE*  $\epsilon$ 4 allele, although this increased A $\beta$  was not associated with any greater impairment in cognition or neuronal function. Importantly, the increase in cortical A $\beta$  in MC  $\epsilon$ 4 carriers was not affected by the *BDNF* Met allele. Thus, allelic variation in *BDNF* and *APOE* may affect different AD processes with  $\epsilon$ 4 increasing cortical A $\beta$  accumulation and *BDNF* Met<sub>66</sub> moderating A $\beta$  related impairment in cognition and neuronal function through its effects on tau.

Neuronal and synaptic loss characteristic of both sporadic and autosomal dominant AD is due to the combined accumulation of A $\beta$  plaques and tau aggregation (Ballatore *et al.* , 2007, Ittner and Götz, 2011, Spires-Jones and Hyman, 2014).

Neuropathological and CSF biomarker studies show that in AD, cognitive impairment and synaptic loss are associated more strongly with the presence and number of neurofibrillary tangles than A $\beta$  plaques (Bennett *et al.* , 2004, Giannakopoulos *et al.* , 2003, Ingelsson *et al.* , 2004). However, neuroimaging studies in preclinical AD report that higher cortical A $\beta$  load is associated with greater rates of cognitive decline and progression to MCI (Lim *et al.* , 2014, Rowe *et al.* , 2013), with these effects mediated by the effect of A $\beta$  on neurodegeneration (Jack and Holtzman, 2013, Lim *et al.* , 2015). In this context, dissociation of the effects of *BDNF* on A $\beta$  and tau associated cognitive impairment observed here are important because they provide evidence that *BDNF* Met<sub>66</sub> influences disease progression through effects on neuronal dysfunction and cognitive impairment associated with tau.

The current observation that *BDNF* Met<sub>66</sub> in preclinical ADAD was associated with increased tau, hippocampal dysfunction and memory impairment is consistent with the role that CNS BDNF plays in synaptic excitation, long-term potentiation and neuronal plasticity (Fahnestock, 2011, Forlenza *et al.* , 2010, Garzon and Fahnestock, 2007, Hariri *et al.* , 2003, Lee *et al.* , 2012, Lu *et al.* , 2013, Peng *et al.* , 2005). Evidence of a mechanistic relationship between BDNF and tau has been shown in cellular studies which demonstrate that BDNF can induce rapid dephosphorylation of tau through TrkB activation (Elliott *et al.* , 2005) and that BDNF loss in AD is specific to tangle-bearing neurons (Ferrer *et al.* , 1999). This has prompted the hypothesis that there may be a direct relationship between CNS BDNF levels and tau (Belrose *et al.* , 2013), although this remains under investigation. Even in the absence of a direct mechanistic link, the large and clinically important effects of *BDNF* Met<sub>66</sub> on memory, hippocampal function and tau, observed in the current ADAD sample, indicate that studying allelic variation in *BDNF* Val66Met may help clarify pathological models of AD and may even provide a reference for the investigation of the effects and clinical consequences of other neurotrophic factors in AD.

As we have noted (Lim *et al.* , 2013), genome-wide association studies (GWAS) of AD do not identify the *BDNF* Val66Met polymorphism as increasing the risk for AD (Lambert *et al.* , 2013). One possible explanation for this is that GWAS typically use a clinical classification of dementia as the target phenotype. Consequently, they may overlook the contribution of *BDNF* because the effects of this gene manifest only in the

earliest stages of the disease (Feng *et al.* , 2013, Lim *et al.* , 2014, Lim *et al.* , 2013). This hypothesis is supported by GWAS of cognitive aging in non-demented older adults, where *BDNF* Val66Met has been associated with memory impairment and decline (Harris and Deary, 2011, Papenberg *et al.* , 2015). Thus, the hypothesis arising from the current and previous studies (Lim *et al.* , 2014, Lim *et al.* , 2013, Lim *et al.* , 2014) is that in studies of cognitive aging, memory decline associated with *BDNF* Met<sub>66</sub> may reflect occult AD as opposed to the effects of normal aging. In contrast to *BDNF*, GWAS of AD identify carriage of *APOE*  $\epsilon$ 4 as increasing risk for AD (Lambert *et al.* , 2013). We have also reported that in preclinical sporadic AD, the *APOE*  $\epsilon$ 4 allele increases the rate of memory decline and brain volume loss associated with high A $\beta$  (Dore *et al.* , 2013, Lim *et al.* , 2014, Lim *et al.* , 2015). We have also observed that A $\beta$ + older adults who carry both the *APOE*  $\epsilon$ 4 and *BDNF* Met<sub>66</sub> allele show greater memory decline than those who carry either one by itself (Lim *et al.* , 2014). Re-analysis of the current data taking into account *APOE*  $\epsilon$ 4 did not indicate any effect of *APOE* or any interaction between *APOE* and *BDNF* on cognition (Table 3). The absence of any effect of *APOE* on cognition in this study is consistent with the results of a detailed meta-analysis of three ADAD cohorts which showed that *APOE* did not moderate age of clinical symptom onset (Ryman *et al.* , 2014). However, despite having no effect on cognitive function or clinical symptom onset, *APOE*  $\epsilon$ 4 was associated with increasing cortical A $\beta$  levels in preclinical MCs. Consequently, one hypothesis for the absence of any *APOE* effect on cognitive and clinical outcomes in ADAD is that these outcomes are related more strongly to neuronal dysfunction and tau than to A $\beta$  accumulation.

This study demonstrates that the deleterious effects of A $\beta$  in ADAD were increased in preclinical individuals who carried the *BDNF* Met<sub>66</sub> allele. Therefore, the results of this study also confirm the similarity between the development of dementia in ADAD and sporadic AD. However, as the current findings are based on cross-sectional data, it will be necessary to replicate these results prospectively. Nonetheless, the strength and consistency of our results with that in sporadic AD is important because they suggest that strategies designed to increase CNS BDNF levels may be a viable therapeutic alternative or addition to those which seek to reduce the neurotoxic effects of A $\beta$ . Our results also suggest strongly that the *BDNF* Val66Met polymorphism should be considered as a potential moderator of clinical trial outcomes in current treatment

and prevention trials in ADAD and sporadic AD (Donohue *et al.* , 2014, Mills *et al.* , 2013).

### **Acknowledgements**

Data collection and sharing for this project was supported by The Dominantly Inherited Alzheimer Network (DIAN; U19AG032438, P50 AG-16570 and P50 AG-005142) funded by the National Institute on Aging (NIA) and the German Center for Neurodegenerative Diseases (DZNE). This work was also supported by the NIHR Queen Square Dementia Biomedical Research Unit and the MRC Dementias Platform UK (MR/L023784/1 and MR/009076/1). We thank the Genome Technology Access Center in the Department of Genetics at Washington University School of Medicine for help with genomic analysis. The Center is partially supported by NCI Cancer Center Support Grant #P30 CA91842 to the Siteman Cancer Center and by ICTS/CTSA Grant# UL1TR000448 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. Data and/or tissue generated from DIAN fibroblasts (or iPSCs) and/or exome chip sequencing was supported by the DIAN-TU Pharma Consortium, (the DIAN-TU Pharma Consortium, <http://dian-tu.wustl.edu/en/pharma-consortium-members/>). This publication is solely the responsibility of the authors and does not necessarily represent the official view of NCRR or NIH. JP Chhatwal is supported by K23-AG049087. YY Lim is supported by the Alzheimer's Australia Dementia Research Fellowship, the Yulgilbar Foundation Postdoctoral Award, and the NHMRC-ARC Dementia Research Development Fellowship (APP1111603).

We thank all participants and their families for their commitment and dedication to helping advance research into the early detection and causation of AD; and the DIAN research and support staff at each of the participating sites for their contributions to this study.

### **Author Contributions**

Study concept and design: YY Lim, P Maruff, PJ Snyder, RJ Bateman

Drafting the manuscript: YY Lim, J Hassenstab, P Maruff, RJ Bateman

Statistical analysis and acquisition of data: YY Lim, J Hassenstab, C Cruchaga, A Goate, AM Fagan, TLS Benzinger, P Maruff, CL Masters, R Allegri, J Chhatwal, MR Farlow, NR Graff-Radford, C Laske, J Levin, E McDade, JM Ringman, M Rossor, S Salloway, PR Schofield, DM Holtzman, JC Morris, RJ Bateman

Interpretation of data: YY Lim, J Hassenstab, P Maruff, RJ Bateman

Revising the manuscript: YY Lim, J Hassenstab, C Cruchaga, A Goate, AM Fagan, TLS Benzinger, P Maruff, PJ Snyder, CL Masters, R Allegri, J Chhatwal, MR Farlow, NR Graff-Radford, C Laske, J Levin, E McDade, JM Ringman, M Rossor, S Salloway, PR Schofield, DM Holtzman, JC Morris, RJ Bateman

### **Conflicts of Interests**

YY Lim, J Hassenstab, C Cruchaga, P Maruff, PJ Snyder, R Allegri, MR Farlow, NR Graff-Radford, C Laske, E McDade, and J Ringman have no disclosures.

A Goate reports grants from NIH, during the conduct of the study and personal fees from Denali Therapeutics, outside the submitted work. AM Fagan reports grants from NIH/NIA, during the conduct of the study, and personal fees from Roche, IBL International, and DiamiR, outside the submitted work. TLS Benzinger reports grants from NIH, during the conduct of the study, and grants from NIH, Avid Radiopharmaceuticals (Eli Lilly), Eli Lilly, and Roche outside the submitted work. CL Masters is a consultant to Prana Biotechnology, Eli Lilly and Actinogen. J Chhatwal reports grants from BrightFocus Foundation, National Institute on Aging, and American Brain Foundation outside the submitted work. J Levin reports grants from DZNE (German Center for Neurodegenerative Diseases), during the conduct of the study; personal fees from Bayer Vital, General Electric, and Willi Gross Foundation; grants from Verum Foundation, ParkinsonFonds Deutschland gGmbH, Deutsche Parkinson Vereinigung, Deutsche Stiftung Neurologie, and Golser Foundation outside the submitted work. M Rossor reports grants from the National Institute on Aging during the conduct of the study. S Salloway reports grants and personal fees from Biogen, Merck, Eli Lilly, Roche and Genentech; and grants from Avid and Functional Neuromodulation outside the submitted work. PR Schofield reports grants from



National Institute on Aging and Mason & Wicking Trusts during the conduct of the study. DM Holtzman reports personal fees from C2N Diagnostics, Genentech, Astra Zeneca, Eli Lilly, Denali Therapeutics and AbbVie outside the submitted work. DM Holtzman also co-founded C2N Diagnostics LLC and have equity interests in the company. JC Morris reports grants from National Institute of Health (P50AG005681, P01AG003991, P01AG026276, UF01AG032438) during the conduct of the study; and consultant fees from Lilly USA and Takeda Pharmaceuticals outside the submitted work. RJ Bateman reports grants from National Institute of Health/National Institute of Aging (U19 AG032438) and Anonymous Foundation during the conduct of the study. RJ Bateman also reports grants from Alzheimer's Association, American Academy of Neurology, Anonymous Foundation, AstraZeneca, BrightFocus Foundation, Cure Alzheimer's Fund, Glenn Foundation for Medical Research, Merck, Metropolitan Life Foundation, National Institute of Health, National Institute of Aging, Pharma Consortium (Biogen Idec, Elan Pharmaceuticals Inc., Eli Lilly and Co., Hoffman La-Roche Inc., Genentech Inc., Janssen Alzheimer Immunotherapy, Mithridion Inc., Novartis Pharma AG, Pfizer Biotherapeutics R and D, Sanofi-Aventi, Eisai), Roche and Ruth K. Broadman Biomedical Research Foundation outside the submitted work. RJ Bateman reports non-financial support from Avid Radiopharmaceuticals; is co-founder of C2N Diagnostics; personal fees and non-financial support from Roche, IMI, Sanofi, Global Alzheimer's Platform, FORUM, OECD, Boehringer Ingelheim; and personal fees from Merck outside the submitted work.

Table 1. Demographic and clinical characteristics

	NC (n=131)	MC Val <sub>66</sub> /Val <sub>66</sub> (n=95)	MC Met <sub>66</sub> (n=48)	<i>p</i>
N (%) Female	58 (44.3%)	37 (38.9%)	25 (52.1%)	.325
N (%) <i>APOE</i> ε4 carrier	39 (29.8%)	24 (25.3%)	12 (25.0%)	.784
Age	<b>38.37 (10.13)</b>	<b>34.45 (8.54)</b>	<b>35.12 (9.51)</b>	<b>.012</b>
Est. Year of Onset	N/A	-12.44 (8.11)	-12.70 (7.60)	.855
Years of Education	14.79 (2.64)	14.72 (3.54)	14.24 (2.56)	.328
GDS	1.24 (1.66)	1.45 (1.83)	1.47 (1.60)	.566
CDR sum of boxes	<b>0.01 (0.06)</b>	<b>0.02 (0.10)</b>	<b>0.06 (0.17)</b>	<b>.005</b>
MMSE	29.20 (1.17)	28.97 (1.37)	29.04 (0.99)	.340

Note: NC = Mutation non-carrier; MC = mutation carrier; GDS = Geriatric Depression Scale; HADS = Hospital Anxiety and Depression Scale; MACQ = Memory Complaints Questionnaire; CDR = Clinical Dementia Rating scale; MMSE = Mini-Mental State Examination

Table 2. Differences in each cognitive marker and biomarker between mutation non-carriers, mutation carriers who are *BDNF* Val<sub>66</sub> homozygotes, and mutation carriers who are *BDNF* Met<sub>66</sub> carriers.

Primary Outcomes	EYO		Group		NC		MC Val <sub>66</sub> /Val <sub>66</sub>		MC Met <sub>66</sub>	
	(df) F	<i>p</i>	(df) F	<i>p</i>	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N
Episodic Memory	<b>(1,268) 20.87</b>	<b>.00</b>	<b>(2,268) 5.35</b>	<b>.00</b>	0.03 (0.82)	131	-0.12 (0.83)	95	-0.43 (0.83)	48
PiB-PET SUVR	<b>(1,223) 18.31</b>	<b>.00</b>	<b>(2,223) 38.85</b>	<b>.00</b>	1.04 (0.54)	106	1.62 (0.53)	82	1.74 (0.53)	39
CSF tau	<b>(1,216) 16.20</b>	<b>.00</b>	<b>(2,216) 19.94</b>	<b>.00</b>	57.18 (40.42)	101	82.83 (40.29)	80	102.15 (40.29)	39
FDG-PET hippocampus	<b>(1,225) 12.13</b>	<b>.00</b>	<b>(2,225) 3.91</b>	<b>.02</b>	1.25 (0.09)	109	1.26 (0.09)	80	1.21 (0.09)	40
Exploratory Outcomes										
Executive Function	(1,268) 2.23	.14	(2,268) 2.10	.12	0.02 (0.80)	131	-0.15 (0.81)	95	-0.22 (0.81)	48
Language	(1,267) 0.28	.60	(2,267) 2.90	.06	0.03 (0.86)	131	-0.15 (0.86)	95	-0.29 (0.86)	48
Attention	(1,268) 2.20	.14	<b>(2,268) 4.00</b>	<b>.02</b>	0.01 (0.81)	131	-0.15 (0.81)	95	-0.37 (0.81)	48
Global Cognition	<b>(1,267) 8.36</b>	<b>.00</b>	<b>(2,267) 3.58</b>	<b>.03</b>	0.02 (0.65)	131	-0.12 (0.65)	95	-0.26 (0.66)	48
CSF Aβ <sub>42</sub>	<b>(1,213) 8.03</b>	<b>.01</b>	<b>(2,213) 7.55</b>	<b>.00</b>	430.72 (147.29)	99	355.78 (146.91)	78	346.57 (146.79)	40
CSF p-tau <sub>181</sub>	<b>(1,217) 8.12</b>	<b>.01</b>	<b>(2,217) 32.22</b>	<b>.00</b>	29.27 (22.75)	101	48.48 (22.69)	80	60.62 (22.67)	40
FDG-PET precuneus	<b>(1,225) 4.62</b>	<b>.03</b>	(2,225) 1.13	.33	2.79 (0.29)	109	2.74 (0.29)	80	2.73 (0.29)	40

Note: EYO = estimated year of symptom onset; Group = effect of group membership as NC, MC Val<sub>66</sub> homozygote or MC Met<sub>66</sub> carrier; all models have been adjusted for estimated year of symptom onset; bolded values are significant at the  $p < .05$  or  $p < .001$  level

Table 3. Effect of estimated year of symptom onset (EYO), *APOE*  $\epsilon$ 4 status, *BDNF* Val66Met status, and the interaction between *APOE* and *BDNF* on each cognitive and biomarker outcome measure

Primary Outcomes	EYO		<i>APOE</i> Group		<i>BDNF</i> Group		<i>APOE</i> x <i>BDNF</i> Group	
	(df) F	<i>p</i>	(df) F	<i>p</i>	(df) F	<i>p</i>	(df) F	<i>p</i>
Episodic Memory	<b>(1,266) 19.32</b>	<b>.00</b>	(1,266) 1.79	.18	<b>(1,266) 3.84</b>	<b>.05</b>	(1,266) 0.08	.77
PiB-PET SUVR	<b>(1,221) 16.73</b>	<b>.00</b>	<b>(1,221) 7.21</b>	<b>.01</b>	(1,221) 2.65	.11	(1,221) 1.18	.28
CSF tau	<b>(1,214) 15.50</b>	<b>.00</b>	(1,214) 0.24	.63	<b>(1,214) 4.06</b>	<b>.04</b>	(1,214) 0.05	.82
FDG-PET hippocampus	<b>(1,223) 11.88</b>	<b>.00</b>	(1,223) 0.05	.83	<b>(1,223) 6.03</b>	<b>.02</b>	(1,223) 0.09	.77
Exploratory Outcomes								
Executive Function	(1,266) 2.50	.12	(1,266) 0.93	.34	(1,266) 0.08	.78	(1,266) 0.08	.78
Language	(1,266) 0.20	.65	(1,266) 0.39	.54	(1,266) 0.99	.32	(1,266) 0.15	.70
Attention	(1,266) 2.40	.12	(1,266) 1.46	.23	(1,266) 2.78	.10	(1,266) 0.49	.49
DIAN Composite	<b>(1,266) 8.14</b>	<b>.01</b>	(1,266) 0.04	.84	(1,266) 0.84	.36	(1,266) 0.06	.81
CSF A $\beta$ <sub>42</sub>	<b>(1,211) 6.94</b>	<b>.01</b>	<b>(1,211) 9.28</b>	<b>.00</b>	(1,211) 0.02	.90	(1,211) 1.39	.24
CSF p-tau <sub>181</sub>	<b>(1,215) 7.39</b>	<b>.01</b>	(1,215) 2.00	.16	<b>(1,215) 4.60</b>	<b>.03</b>	(1,215) 0.34	.56
FDG-PET precuneus	<b>(1,223) 4.82</b>	<b>.03</b>	(1,223) 0.65	.42	(1,223) 0.001	.98	(1,223) 0.02	.91

Note: all models have been adjusted for estimated year of symptom onset (EYO); *APOE* Group indicates effect of group membership as NC, *APOE*  $\epsilon$ 4 carrier or *APOE*  $\epsilon$ 4 non-carrier; *BDNF* Group indicates effect of group membership as NC, MC Val<sub>66</sub> homozygote or MC Met<sub>66</sub> carrier; bolded values are significant at the  $p < .05$  level

Table 4. Re-analysis of the effect of *BDNF* Val66Met on each primary outcome variable, covarying for the potential confounding effect of sex


	EYO		Sex		<i>BDNF</i> Group	
	(df) F	<i>p</i>	(df) F	<i>p</i>	(df) F	<i>p</i>
Episodic Memory	<b>(1,267) 13.96</b>	<b>.00</b>	<b>(1,267) 8.24</b>	<b>.00</b>	<b>(2,267) 5.17</b>	<b>.00</b>
PiB-PET SUVR	<b>(1,222) 18.43</b>	<b>.00</b>	(1,222) 0.01	.92	<b>(2,222) 40.59</b>	<b>.00</b>
CSF tau	<b>(1,215) 17.02</b>	<b>.00</b>	(1,215) 0.56	.46	<b>(2,215) 20.98</b>	<b>.00</b>
FDG-PET hippocampus	<b>(1,224) 5.60</b>	<b>.02</b>	(1,224) 1.35	.25	<b>(2,224) 3.95</b>	<b>.02</b>


### Figure Captions

Figure 1. Magnitude of cognitive impairment in preclinical MC Val<sub>66</sub> homozygotes and preclinical MC Met<sub>66</sub> carriers when compared to mutation non-carriers (error bars represent 95% confidence intervals) (statistical significance occurs when 95% confidence intervals do not cross “0” line).


Figure 2. Magnitude of abnormality on markers of A $\beta$ , tau and glucose metabolism in preclinical MC Val<sub>66</sub> homozygotes and preclinical MC Met<sub>66</sub> carriers when compared to mutation non-carriers (error bars represent 95% confidence intervals) (statistical significance occurs when 95% confidence intervals do not cross “0” line).

Figure 3. Relationship between estimated year of clinical symptom onset and episodic memory performance (A), glucose metabolism in the hippocampus (B), cortical A $\beta$  levels (C), and CSF p-tau<sub>181</sub> levels (D), in mutation non-carriers, preclinical MC Val<sub>66</sub> homozygotes, and preclinical MC Met<sub>66</sub> carriers.









## References

- Ballatore C, Lee VMY, Trojanowski JQ. Tau-mediated neurodegeneration in Alzheimer's disease and related disorders. *Nature Reviews, Neuroscience*. 2007;8:663-72.
- Bateman RJ, Xiong C, Benzinger TLS, Fagan AM, Goate A, Fox NC, et al. Clinical and biomarker changes in Dominantly Inherited Alzheimer's disease. *The New England Journal of Medicine*. 2012;367(9):795-804.
- Belrose JC, Masoudi R, Michalski B, Fahnstock M. Increased pro-nerve growth factor and decreased brain-derived neurotrophic factor in non-Alzheimer's disease tauopathies. *Neurobiology of Aging*. 2013;35(4):926-33.
- Bennett DA, Schneider JA, Wilson RS, Bienias JL, Arnold SE. Neurofibrillary tangles mediate the association of amyloid load with clinical Alzheimer disease and level of cognitive function. *Archives of Neurology*. 2004;61(3):378-84.
- Benzinger TL, Blazey T, Jack CR, Koeppe RA, Su Y, Xiong C, et al. Regional variability of imaging biomarkers in autosomal dominant Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;110(47):E4502-E9.
- Buchman AS, Yu L, Boyle PA, Schneider JA, De Jager PL, Bennett DA. Higher brain BDNF gene expression is associated with slower cognitive decline in older adults. *Neurology*. 2016;86(8):735-41.
- Caccamo A, Maldonado MA, Bokov AF, Majumder S, Oddo S. CBP gene transfer increases BDNF levels and ameliorates learning and memory deficits in a mouse model of Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;107(52):22687-92.

- Chen ZY, Jing D, Bath KG, Ieraci A, Khan T, Siao CJ, et al. Genetic variant BDNF (Val66Met) polymorphism alters anxiety-related behavior. *Science*. 2006;314(5796):140-3.
- Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. New Jersey: Lawrence Erlbaum Associates; 1988.
- Donohue MC, Sperling RA, Salmon DP, Rentz DM, Raman R, Thomas RG, et al. The Preclinical Alzheimer Cognitive Composite: Measuring amyloid-related decline. *JAMA Neurology*. 2014;71(8):961-70.
- Dore V, Villemagne VL, Bourgeat P, Fripp J, Acosta O, Chetelat G, et al. Cross-sectional and longitudinal analysis of the relationship between A $\beta$  deposition, cortical thickness, and memory in cognitively unimpaired individuals and in Alzheimer's disease. *JAMA Neurology*. 2013;70(7):903-11.
- Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, et al. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell*. 2003;112:257-69.
- Elliott E, Atlas R, Lange A, Ginzburg I. Brain-derived neurotrophic factor induces a rapid dephosphorylation of tau protein through a PI-3 kinase signalling mechanism. *European Journal of Neuroscience*. 2005;22:1081-9.
- Fahnestock M. Brain-derived neurotrophic factor: The link between amyloid-b and memory loss. *Future Neurology*. 2011;6:627-39.
- Feng S, Sevigny J, Verma A, Bennett D, Lim YY, Maruff P. Genetic and image biomarkers associated with decline in cognitive measures and brain glucose metabolism in populations of early Alzheimer's disease. *Alzheimer's & Dementia*. 2013;9:178.
- Ferrer I, Marín C, Rey MJ, Ribalta T, Goutan E, Blanco RT, et al. BDNF and full-length and truncated TrkB expression in Alzheimer disease: Implications in therapeutic

strategies. *Journal of Neuropathology and Experimental Neurology*.

1999;58(7):729-39.

Forlenza OV, Diniz BS, Teixeira AL, Ojopi EB, Talib LL, Mendonça VA, et al. Effect of brain-derived neurotrophic factor Val66Met polymorphism and serum levels on the progression of mild cognitive impairment. *World Journal of Biological Psychiatry*. 2010;11(6):774-80.

Garzon DJ, Fahnestock M. Oligomeric amyloid decreases basal levels of brain-derived neurotrophic factor (BDNF) mRNA via specific downregulation of BDNF transcripts IV and V in differentiated human neuroblastoma cells. *The Journal of Neuroscience*. 2007;27(10):2628-35.

Giannakopoulos P, Herrmann FR, Bussière T, Bouras C, Kövari E, Perl DP, et al. Tangle and neuron numbers, but not amyloid load, predict cognitive status in Alzheimer's disease. *Neurology*. 2003;60(9):1495-500.

Hardy JA, Higgins GA. Alzheimer's disease: The amyloid cascade hypothesis. *Science*. 1992;256:184-5.

Hariri AR, Goldberg TE, Mattay VS, Kolachana BS, Callicott JH, Egan MF, et al. Brain-derived neurotrophic factor val66met polymorphism affects human memory-related hippocampal activity and predicts memory performance. *The Journal of Neuroscience*. 2003;23(17):6690-4.

Harris SE, Deary IJ. The genetics of cognitive ability and cognitive ageing in healthy older people. *Trends in Cognitive Sciences*. 2011;15(9):388-94.

Ingelsson M, Fukumoto H, Newell KL, Growdon JH, Hedley-Whyte ET, Frosch MP, et al. Early Abeta accumulation and progressive synaptic loss, gliosis, and tangle formation in AD brain. *Neurology*. 2004;62:925-31.

- Ittner LM, Götz J. Amyloid- $\beta$  and tau - a toxic pas de deux in Alzheimer's disease. *Nature Reviews, Neuroscience*. 2011;12(2):65-72.
- Jack CR, Holtzman DM. Biomarker modeling of Alzheimer's disease. *Neuron*. 2013;80(6):1347-58.
- Keaney JFJ, Larson MG, Vasan RS, Wilson PW, Lipinska I, Corey D, et al. Obesity and systemic oxidative stress: Clinical correlates of oxidative stress in the Framingham Study. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2003;23(3):434-9.
- Kim A, Fagan AM, Goate AM, Benzinger TL, Morris JC, Head D, et al. Lack of an association of BDNF Val66Met polymorphism and plasma BDNF with hippocampal volume and memory. *Cognitive, Affective & Behavioral Neuroscience*. 2015;15(3):625-43.
- Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C, et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nature Genetics*. 2013;45(12):1452-8.
- Lee ST, Chu K, Jung KH, Kim JH, Huh JY, Yoon H, et al. miR-206 regulates brain-derived neurotrophic factor in Alzheimer disease model. *Annals of Neurology*. 2012;72:269-77.
- Lim YY, Maruff P, Pietrzak RH, Ames D, Ellis KA, Harrington K, et al. Effect of amyloid on memory and non-memory decline from preclinical to clinical Alzheimer's disease. *Brain*. 2014;137:221-31.
- Lim YY, Pietrzak RH, Bourgeat P, Ames D, Ellis KA, Rembach A, et al. Relationships between performance on the Cogstate Brief Battery, neurodegeneration, and A $\beta$  accumulation in cognitively normal older adults and adults with MCI. *Archives of Clinical Neuropsychology*. 2015;30(1):49-58.

- Lim YY, Villemagne VL, Laws SM, Ames D, Pietrzak RH, Ellis KA, et al. BDNF Val66Met moderates A $\beta$ -related memory decline and hippocampal atrophy in prodromal Alzheimer's disease: A preliminary study. *PLoS One*. 2014;9:e86498.
- Lim YY, Villemagne VL, Laws SM, Ames D, Pietrzak RH, Ellis KA, et al. BDNF Val66Met, A $\beta$  amyloid and cognitive decline in preclinical Alzheimer's disease. *Neurobiology of Aging*. 2013;34:2457-64.
- Lim YY, Villemagne VL, Laws SM, Pietrzak RH, Snyder PJ, Ames D, et al. APOE and BDNF polymorphisms moderate amyloid  $\beta$ -related cognitive decline in preclinical Alzheimer's disease. *Molecular Psychiatry*. 2014;epub.
- Lim YY, Villemagne VL, Pietrzak RH, Ames D, Ellis KA, Harrington K, et al. APOE  $\epsilon$ 4 moderates amyloid-related memory decline in preclinical Alzheimer's disease. *Neurobiology of Aging*. 2015;36(3):1239-44.
- Lu B, Nagappan G, Guan X, Nathan PJ, Wren P. BDNF-based synaptic repair as a disease-modifying strategy for neurodegenerative diseases. *Nature Reviews, Neuroscience*. 2013;14:401-16.
- Mills SM, Mallmann J, Santacruz AM, Fuqua A, Carril M, Aisen PS, et al. Preclinical trials in autosomal dominant AD: Implementation of the DIAN-TU trial. *Revue Neurologique*. 2013;169:737-43.
- Morris JC. The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology*. 1983;43:2412-4.
- Morris JC, Aisen PS, Bateman RJ, Benzinger T, Cairns NJ, Fagan AM, et al. Developing an international network for Alzheimer's research: The Dominantly Inherited Alzheimer Network. *Clinical Investigation*. 2012;2(10):975-84.

- Papenberg G, Salami A, Persson J, Lindenberger U, Bäckman L. Genetics and functional Imaging: Effects of APOE, BDNF, COMT, and KIBRA in aging. *Neuropsychology Review*. 2015;25(1):47-62.
- Peng S, Wu J, Mufson EJ, Fahnstock M. Precursor form of brain-derived neurotrophic factor and mature brain-derived neurotrophic factor are decreased in the preclinical stages of Alzheimer's disease. *Journal of Neurochemistry*. 2005;93:1412-21.
- Rosa E, Fahnstock M. CREB expression mediates amyloid  $\beta$ -induced basal BDNF downregulation. *Neurobiology of Aging*. 2015;36:2406-13.
- Rowe CC, Bourgeat P, Ellis KA, Brown B, Lim YY, Mulligan R, et al. Predicting Alzheimer disease with  $\beta$ -amyloid imaging: Results from the Australian imaging, biomarkers, and lifestyle study of ageing. *Annals of Neurology*. 2013;74(6):905-13.
- Ryman DC, Acosta-Baena N, Aisen PS, Bird T, Danek A, Fox NC, et al. Symptom onset in autosomal dominant Alzheimer disease: A systematic review and meta-analysis. *Neurology*. 2014;85(3):253-60.
- Spires-Jones T, Hyman BT. The intersection of amyloid beta and tau at synapses in Alzheimer's disease. *Neuron*. 2014;82:756-71.
- Storandt M, Balota DA, Aschenbrenner AJ, Morris JC. Clinical and psychological characteristics of the initial cohort of the Dominantly Inherited Alzheimer Network (DIAN). *Neuropsychology*. 2014;28(1):19-29.
- Su Y, Blazey TM, Snyder AZ, Raichle ME, Marcus DS, Ances BM, et al. Partial volume correction in quantitative amyloid imaging. *NeuroImage*. 2015;107:55-64.
- Talbot C, Lendon C, Craddock N, Shears S, Morris JC, Goate A. Protection against Alzheimer's disease with apoE epsilon 2. *Lancet*. 1994;343(8910):1432-3.