Appendix M. Included Studies

Abernethy AP, Currow DC, Frith P, Fazekas BS, McHugh A, and Bui C. Randomised, double blind, placebo controlled crossover trial of sustained release morphine for the management of refractory dyspnoea. BMJ 2003;327(7414):523-8. [Rec#: 50928]
INCLUDE: considered for evidence table

Abratt R and Viljoen G. Assessment of quality of life by clinicians—experience of a practical method in lung cancer patients. S Afr Med J 95;85(9):896-8. [Rec#: 30022]
INCLUDE: considered for evidence table

Addington-Hall J, Fakhoury W, and McCarthy M. Specialist palliative care in nonmalignant disease. Palliat Med 98;12(6):417-27. [Rec#: 9055]
INCLUDE: considered for evidence table

Addington-Hall JM, MacDonald LD, Anderson HR, Chamberlain J, Freeling P, Bland JM, and Raftery J. Randomised controlled trial of effects of coordinating care for terminally ill cancer patients. BMJ 92;305(6865):1317-22. [Rec#: 9059]
INCLUDE: considered for evidence table

Addington-Hall, J, Altmann, D, and McCarthy, M. Which terminally ill cancer patients receive hospice in-patient care? Social Science & Medicine 98;46(8):1011-6. [Rec#: 50168]
INCLUDE: considered for evidence table

Addington-Hall, J, Lay, M, Altmann, D, and McCarthy, M. Symptom control, communication with health professionals, and hospital care of stroke patients in the last year of life as reported by surviving family, friends, and officials. Stroke 95;26(12):2242-8. [Rec#: 50169]
INCLUDE: considered for evidence table

Ahrens T, Yancey V, and Kollef M. Improving family communications at the end of life: implications for length of stay in the intensive care unit and resource use. Am J Crit Care 2003;12(4):317-23; discussion 324. [Rec#: 77]
INCLUDE: considered for evidence table

Ahronheim JC, Morrison RS, Baskin SA, Morris J, and Meier DE. Treatment of the dying in the acute care hospital. Advanced dementia and metastatic cancer. Arch Intern Med 96;156(18):2094-100. [Rec#: 79]
INCLUDE: considered for evidence table

Albanese MC, Plewka M, Gregori D, Fresco C, Avon G, Caliandro D, Grassi L, Rossi P, Udina P, Bulfoni A, Colle R, Cuzzato AL, and Fioretti PM. Use of medical resources and quality of life of patients with chronic heart failure: a prospective survey in a large Italian community hospital. Eur J Heart Fail 99;1(4):411-7. [Rec#: 30090]
INCLUDE: considered for evidence table

Albert SM, Murphy PL, Del Bene ML, and Rowland LP. Prospective study of palliative care in ALS: choice, timing, outcomes. J Neurol Sci 99;169(1-2):108-13. [Rec#: 103]
INCLUDE: considered for evidence table

Allen, RS, DeLaine, SR, Chaplin, WF, Marson, DC, Bourgeois, MS, Dijkstra, K, and Burgio, LD. Advance care planning in nursing homes: correlates of capacity and possession of advance directives. Gerontologist 2003;43(3):309-17. [Rec#: 50173]
INCLUDE: considered for evidence table

Allen, RS, Haley, WE, Small, BJ, and McMillan, SC. Pain reports by older hospice cancer patients and family caregivers: the role of cognitive functioning. Gerontologist 2002;42(4):507-14. [Rec#: 50174]
INCLUDE: considered for evidence table

Anderson H, Ward C, Eardley A, Gomm SA, Connolly M, Coppinger T, Corgie D, Williams JL, and Makin WP. The concerns of patients under palliative care and a heart failure clinic are not being met. Palliat Med 2001;15(4):279-86. [Rec#: 183]
INCLUDE: considered for evidence table

Anderson JP, Kaplan RM, and Schneiderman LJ. Effects of offering advance directives on quality adjusted life expectancy and psychological well-being among ill adults. J Clin Epidemiol 94;47(7):761-72. [Rec#: 186]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Andrews SC. Caregiver burden and symptom distress in people with cancer receiving hospice care. Oncol Nurs Forum 2001;28(9):1469-74. [Rec#: 196]
INCLUDE: considered for evidence table

Ankrom M, Zelesnick L, Barofsky I, Georas S, Finucane TE, and Greenough WB 3rd. Elective discontinuation of life-sustaining mechanical ventilation on a chronic ventilator unit. J Am Geriatr Soc 2001;49(11):1549-54. [Rec#: 202]
INCLUDE: considered for evidence table

Aranda SK and Hayman-White K. Home caregivers of the person with advanced cancer: an Australian perspective. Cancer Nurs 2001;24(4):300-7. [Rec#: 221]
INCLUDE: considered for evidence table

Arblaster G, Brooks D, Hudson R, and Petty M. Terminally ill patients' expectations of nurses. Aust J Adv Nurs 90;7(3):34-43. [Rec#: 9201]
INCLUDE: considered for evidence table

Armes PJ and Higginson IJ. What constitutes high-quality HIV/AIDS palliative care? J Palliat Care 99;15(4):5-12. [Rec#: 9215]
INCLUDE: considered for evidence table

Asch DA, Faber-Langendoen K, Shea JA, and Christakis NA. The sequence of withdrawing life-sustaining treatment from patients. Am J Med 99;107(2):153-6. [Rec#: 242]
INCLUDE: considered for evidence table

Ashbury FD, Findlay H, Reynolds B, and McKerracher K. A Canadian survey of cancer patients' experiences: are their needs being met? J Pain Symptom Manage 98;16(5):298-306. [Rec#: 245]
INCLUDE: considered for evidence table

Astradsson E, Granath L, Heedman PA, and Starkhammar H. Cancer patients hospitalised for palliative reasons. Symptoms and needs presented at a university hospital. Support Care Cancer 2001;9(2):97-102. [Rec#: 254]
INCLUDE: considered for evidence table

Azoulay, E, Chevret, S, Leleu, G, Pochard, F, Barboteu, M, Adrie, C, Canoui, P, Le Gall, J. R, and Schlemmer, B. Half the families of intensive care unit patients experience inadequate communication with physicians. Critical Care Medicine 2000;28(8):3044-9. [Rec#: 50193]
INCLUDE: considered for evidence table

Baer WM and Hanson LC. Families' perception of the added value of hospice in the nursing home. J Am Geriatr Soc 2000;48(8):879-82. Comment in: J Am Geriatr Soc. 2000 Aug;48(8):1017-8. PMID: 10968312. Comment in: J Am Geriatr Soc. 2001 Apr;49(4):492. PMID: 11347800. [Rec#: 296]
INCLUDE: considered for evidence table

Baines BK and Norlander L. The relationship of pain and suffering in a hospice population. Am J Hosp Palliat Care 2000;17(5):319-26. [Rec#: 309]
INCLUDE: considered for evidence table

Bakas T, Lewis RR, and Parsons JE. Caregiving tasks among family caregivers of patients with lung cancer. Oncol Nurs Forum 2001;28(5):847-54. [Rec#: 30289]
INCLUDE: considered for evidence table

Baker NT and Seager RD. A comparison of the psychosocial needs of hospice patients with AIDS and those with other diagnoses. Hosp J 91;7(1-2):61-9. [Rec#: 323]
INCLUDE: considered for evidence table

Baker R, Wu AW, Teno JM, Kreling B, Damiano AM, Rubin HR, Roach MJ, Wenger NS, Phillips RS, Desbiens NA, Connors AF Jr, Knaus W, and Lynn J. Family satisfaction with end-of-life care in seriously ill hospitalized adults. J Am Geriatr Soc 2000;48(5 Suppl):S61-9. [Rec#: 324]
INCLUDE: considered for evidence table

Balmer CE, Thomas P, and Osborne RJ. Who wants second-line, palliative chemotherapy? Psychooncology 2001;10(5):410-8. [Rec#: 351]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Barry LC, Kasl SV, and Prigerson HG. Psychiatric disorders among bereaved persons: the role of perceived circumstances of death and preparedness for death. Am J Geriatr Psychiatry 2002;10(4):447-57. [Rec#: 394]
INCLUDE: considered for evidence table

Bass DM and Bowman K. The transition from caregiving to bereavement: the relationship of care-related strain and adjustment to death. Gerontologist 90;30(1):35-42. [Rec#: 412]
INCLUDE: considered for evidence table

Bass DM, Bowman K, and Noelker LS. The influence of caregiving and bereavement support on adjusting to an older relative's death. Gerontologist 91;31(1):32-42. [Rec#: 413]
INCLUDE: considered for evidence table

Bauduer F, Capdupuy C, and Renoux M. Characteristics of deaths in a department of oncohaematology within a general hospital. A study of 81 cases. Support Care Cancer 2000;8(4):302-6. [Rec#: 433]
INCLUDE: considered for evidence table

Beck-Friis B and Strang P. The family in hospital-based home care with special reference to terminally ill cancer patients. J Palliat Care 93;9(1):5-13. [Rec#: 447]
INCLUDE: considered for evidence table

Beguin C, France FR, and Ninane J. Systematic analysis of in-patients' circumstances and causes of death: a tool to improve quality of care. Int J Qual Health Care 97;9(6):427-33. [Rec#: 40235]
INCLUDE: considered for evidence table

Bennett SJ, Perkins SM, Lane KA, Deer M, Brater DC, and Murray MD. Social support and health-related quality of life in chronic heart failure patients. Qual Life Res 2001;10(8):671-82. [Rec#: 30411]
INCLUDE: considered for evidence table

Bercovitch M, Waller A, and Adunsky A. High dose morphine use in the hospice setting. A database survey of patient characteristics and effect on life expectancy. Cancer 99;86(5):871-7. Comment in: Cancer. 2000 Jul 1;89(1):220-1. PMID: 10897022. [Rec#: 499]
INCLUDE: considered for evidence table

Bernabei, R, Gambassi, G, Lapane, K, Landi, F, Gatsonis, C, Dunlop, R, Lipsitz, L, Steel, K, and Mor, V. Management of pain in elderly patients with cancer. SAGE Study Group. Systematic Assessment of Geriatric Drug Use via Epidemiology. JAMA 98;279(23):1877-82. [Rec#: 50214]
INCLUDE: considered for evidence table

Blackhall LJ, Frank G, Murphy ST, Michel V, Palmer JM, and Azen SP. Ethnicity and attitudes towards life sustaining technology. Soc Sci Med 99;48(12):1779-89. [Rec#: 591]
INCLUDE: considered for evidence table

Blackhall LJ, Murphy ST, Frank G, Michel V, and Azen S. Ethnicity and attitudes toward patient autonomy. JAMA 95;274(10):820-5. Comment in: JAMA. 1995 Sep 13;274(10):844-5. PMID: 7650810. Comment in: JAMA. 1996 Jan 10;275(2):107; author reply 109-10. PMID: 8531296. Comment in: JAMA. 1996 Jan 10;275(2):107; author reply 109-10. PMID: 8531297. [Rec#: 592]
INCLUDE: considered for evidence table

Bodnar JC and Kiecolt-Glaser JK. Caregiver depression after bereavement: chronic stress isn't over when it's over. Psychol Aging 94;9(3):372-80. [Rec#: 623]
INCLUDE: considered for evidence table

Bookbinder M, Coyle N, Kiss M, Goldstein ML, Holritz K, Thaler H, Gianella A, Derby S, Brown M, Racolin A, Ho MN, and Portenoy RK. Implementing national standards for cancer pain management: program model and evaluation. J Pain Symptom Manage 96;12(6):334-47; discussion 331-3. [Rec#: 659]
INCLUDE: considered for evidence table

Bookwala J, Coppola KM, Fagerlin A, Ditto PH, Danks JH, and Smucker WD. Gender differences in older adults' preferences for life-sustaining medical treatments and end-of-life values. Death Stud 2001;25(2):127-49. [Rec#: 660]
INCLUDE: considered for evidence table

Booth S, Kelly MJ, Cox NP, Adams L, and Guz A. Does oxygen help dyspnea in patients with cancer? Am J Respir Crit Care Med 96;153(5):1515-8. [Rec#: 665]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Borum ML, Lynn J, and Zhong Z. The effects of patient race on outcomes in seriously ill patients in SUPPORT: an overview of economic impact, medical intervention, and end-of-life decisions. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. J Am Geriatr Soc 2000;48(5 Suppl):S194-8. [Rec#: 685]
INCLUDE: considered for evidence table

Bosek MS, Lowry E, Lindeman DA, Burck JR, and Gwyther LP. Promoting a good death for persons with dementia in nursing facilities: family caregivers' perspectives. JONAS Healthc Law Ethics Regul 2003;5(2):34-41. [Rec#: 690]
INCLUDE: considered for evidence table

Bostrom B, Sandh M, Lundberg D, and Fridlund B. A comparison of pain and health-related quality of life between two groups of cancer patients with differing average levels of pain. J Clin Nurs 2003;12(5):726-35. [Rec#: 697]
INCLUDE: considered for evidence table

Bostrom B, Sandh M, Lundberg D, and Fridlund B. Cancer-related pain in palliative care: patients' perceptions of pain management. J Adv Nurs 2004;45(4):410-9. [Rec#: 696]
INCLUDE: considered for evidence table

Boyd KJ and Kelly M. Oral morphine as symptomatic treatment of dyspnoea in patients with advanced cancer. Palliat Med 97;11(4):277-81. [Rec#: 731]
INCLUDE: considered for evidence table

Bradley EH, Hallemeier AG, Fried TR, Johnson-Hurzeler R, Cherlin EJ, Kasl SV, and Horwitz SM. Documentation of discussions about prognosis with terminally ill patients. Am J Med 2001;111(3):218-23. [Rec#: 743]
INCLUDE: considered for evidence table

Bradley EH, Peiris V, and Wetle T. Discussions about end-of-life care in nursing homes. J Am Geriatr Soc 98;46(10):1235-41. [Rec#: 744]
INCLUDE: considered for evidence table

Brazil K, Bedard M, Willison K, and Hode M. Caregiving and its impact on families of the terminally ill. Aging and Mental Health 2003;7(5):376-82. [Rec#: 50230]
INCLUDE: considered for evidence table

Breitbart W, Gibson C, and Tremblay A. The delirium experience: delirium recall and delirium-related distress in hospitalized patients with cancer, their spouses/caregivers, and their nurses. Psychosomatics 2002;43(3):183-94. [Rec#: 50234]
INCLUDE: considered for evidence table

Breitbart W, McDonald MV, Rosenfeld B, Passik SD, Hewitt D, Thaler H, and Portenoy R. K. Pain in ambulatory AIDS patients. I: Pain characteristics and medical correlates. Pain 96;68(2-3):315-21. [Rec#: 50236]
INCLUDE: considered for evidence table

Breitbart W, Tremblay A, and Gibson C. An open trial of olanzapine for the treatment of delirium in hospitalized cancer patients. Psychosomatics 2002;43(3):175-82. [Rec#: 50237]
INCLUDE: considered for evidence table

Brown D, Roberts JA, Elkins TE, Larson D, and Hopkins M. Hard choices: the gynecologic cancer patient's end-of-life preferences. Gynecol Oncol 94;55(3 Pt 1):355-62. [Rec#: 835]
INCLUDE: considered for evidence table

Brown JE, Brown RF, Miller RM, Dunn SM, King MT, Coates AS, and Butow PN. Coping with metastatic melanoma: the last year of life. Psychooncology 2000;9(4):283-92. [Rec#: 844]
INCLUDE: considered for evidence table

Brown R, Butow PN, Boyer M J, and Tattersall M. H. Promoting patient participation in the cancer consultation: evaluation of a prompt sheet and coaching in question-asking. British Journal of Cancer 99;80(1-2):242-8. [Rec#: 50244]
INCLUDE: considered for evidence table

Bruera E, de Stoutz N, Velasco-Leiva A, Schoeller T, and Hanson J. Effects of oxygen on dyspnea in hypoxaemic terminal-cancer patients. Lancet 93;342(8862):13-4. [Rec#: 866]
INCLUDE: considered for evidence table

Bruera E, Kuehn N, Emery B, Macmillan K, and Hanson J. Social and demographic characteristics of patients admitted to a palliative care unit. J Palliat Care 90;6(4):16-20. [Rec#: 867]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Bruera E, MacEachern T, Ripamonti C, and Hanson J. Subcutaneous morphine for dyspnea in cancer patients. Ann Intern Med 93;119(9):906-7. Comment in: Ann Intern Med. 1994 Apr 15;120(8):692-3. PMID: 8192764. [Rec#: 9839]
INCLUDE: considered for evidence table

Bruera E, Miller MJ, Kuehn N, MacEachern T, and Hanson J. Estimate of survival of patients admitted to a palliative care unit: a prospective study. J Pain Symptom Manage 92;7(2):82-6. [Rec#: 870]
INCLUDE: considered for evidence table

Bruera E, Neumann CM, Gagnon B, Brenneis C, Kneisler P, Selmser P, and Hanson J. Edmonton Regional Palliative Care Program: impact on patterns of terminal cancer care. CMAJ 99;161(3):290-3. [Rec#: 9842]
INCLUDE: considered for evidence table

Bruera E, Sloan P, Mount B, Scott J, and Suarez-Almazor M. A randomized, double-blind, double-dummy, crossover trial comparing the safety and efficacy of oral sustained-release hydromorphone with immediate-release hydromorphone in patients with cancer pain. Canadian Palliative Care Clinical Trials Group. J Clin Oncol 96;14(5):1713-7. [Rec#: 877]
INCLUDE: considered for evidence table

Bruera E, Sweeney C, Calder K, Palmer L, and Benisch-Tolley S. Patient preferences versus physician perceptions of treatment decisions in cancer care. J Clin Oncol 2001;19(11):2883-5. [Rec#: 879]
INCLUDE: considered for evidence table

Bruera E, Sweeney C, Willey J, Palmer JL, Strasser F, and Strauch E. Perception of discomfort by relatives and nurses in unresponsive terminally ill patients with cancer: a prospective study. J Pain Symptom Manage 2003;26(3):818-26. [Rec#: 881]
INCLUDE: considered for evidence table

Bruera E, Belzile M, Neumann, CM, Ford I, Harsanyi Z, and Darke A. Twice-daily versus once-daily morphine sulphate controlled-release suppositories for the treatment of cancer pain. A randomized controlled trial. Supportive Care in Cancer 99;7(4):280-3. [Rec#: 50246]
INCLUDE: considered for evidence table

Bruera E, Brenneis C, Michaud M, Bacovsky R, Chadwick S, Emeno A, and MacDonald N. Use of the subcutaneous route for the administration of narcotics in patients with cancer pain. Cancer 88;62(2):407-11. [Rec#: 50249]
INCLUDE: considered for evidence table

Bruera E, Brenneis C, Michaud M, Chadwick S, and MacDonald RN. Continuous sc infusion of narcotics using a portable disposable device in patients with advanced cancer. Cancer Treatment Reports 87;71(6):635-7. [Rec#: 50250]
INCLUDE: considered for evidence table

Bruera E, Carraro S, Roca E, Cedaro L, and Chacon R. Association between malnutrition and caloric intake, emesis, psychological depression, glucose taste, and tumor mass. Cancer Treatment Reports 84;68(6):873-6. [Rec#: 50259]
INCLUDE: considered for evidence table

Bruera E, Macmillan K, Hanson J, and MacDonald RN. The cognitive effects of the administration of narcotic analgesics in patients with cancer pain. Pain 89;39(1):13-6. [Rec#: 50277]
INCLUDE: considered for evidence table

Bruera E, MacMillan K, Hanson J, and MacDonald RN. The Edmonton staging system for cancer pain: preliminary report. Pain 89;37(2):203-9. [Rec#: 50279]
INCLUDE: considered for evidence table

Bruera E, Macmillan K, Kuehn N, and Miller MJ. Circadian distribution of extra doses of narcotic analgesics in patients with cancer pain: a preliminary report. Pain 92;49(3):311-4. [Rec#: 50281]
INCLUDE: considered for evidence table

Bruera E, Michaud M, Vigano A, Neumann CM, Watanabe S, and Hanson J. Multidisciplinary symptom control clinic in a cancer center: a retrospective study. Supportive Care in Cancer 2001;9(3):162-8. [Rec#: 50286]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Bruera E, Miller MJ, Macmillan K, and Kuehn N. Neuropsychological effects of methylphenidate in patients receiving a continuous infusion of narcotics for cancer pain. Pain 92;48(2):163-6. [Rec#: 50288]
INCLUDE: considered for evidence table

Bruera E, Pituskin E, Calder K, Neumann CM, and Hanson J. The addition of an audiocassette recording of a consultation to written recommendations for patients with advanced cancer: A randomized, controlled trial. Cancer 99;86(11):2420-5. [Rec#: 50294]
INCLUDE: considered for evidence table

Bruera E, Watanabe S, Fainsinger RL, Spachynski K, Suarez-Almazor M, and Inturrisi C. Custom-made capsules and suppositories of methadone for patients on high-dose opioids for cancer pain. Pain 95;62(2):141-6. [Rec#: 50305]
INCLUDE: considered for evidence table

Bruera E, Willey JS, Palmer JL, and Rosales M. Treatment decisions for breast carcinoma: patient preferences and physician perceptions. Cancer 2002;94(7):2076-80. [Rec#: 50306]
INCLUDE: considered for evidence table

Brumley RD, Enguidanos S, and Cherin DA. Effectiveness of a home-based palliative care program for end-of-life. J Palliat Med 2003;6(5):715-24. [Rec#: 885]
INCLUDE: considered for evidence table

Buchanan RJ, Choi M, Wang S, and Huang C. Analyses of nursing home residents in hospice care using the minimum data set. Palliat Med 2002;16(6):465-80. [Rec#: 897]
INCLUDE: considered for evidence table

Bucher JA, Loscalzo M, Zabora, J, Houts PS, Hooker C, and BrintzenhofeSzoc K. Problem-solving cancer care education for patients and caregivers. Cancer Practice 2001;9(2):66-70. [Rec#: 50309]
INCLUDE: considered for evidence table

Burge F, Lawson B, and Johnston G. Family physician continuity of care and emergency department use in end-of-life cancer care. Med Care 2003;41(8):992-1001. [Rec#: 917]
INCLUDE: considered for evidence table

Burge F, Lawson B, Johnston G, and Cummings I. Primary care continuity and location of death for those with cancer. J Palliat Med 2003;6(6):911-8. [Rec#: 919]
INCLUDE: considered for evidence table

Burgio L, Stevens A, Guy D, Roth DL, and Haley WE. Impact of two psychosocial interventions on white and African American family caregivers of individuals with dementia. Gerontologist 2003;43(4):568-79. [Rec#: 50915]
INCLUDE: considered for evidence table

Burns CM, Dixon T, Broom D, Smith WT, and Craft PS. Family caregiver knowledge of treatment intent in a longitudinal study of patients with advanced cancer. Support Care Cancer 2003;11(10):629-37. [Rec#: 40484]
INCLUDE: considered for evidence table

Burns R, Nichols LO, Martindale-Adams J, Graney MJ, and Lummus A. Primary care interventions for dementia caregivers: 2-year outcomes from the REACH study. Gerontologist 2003;43(4):547-55. [Rec#: 50916]
INCLUDE: considered for evidence table

Busse R, Wagner HP, Krauth C, Klein-Lange M, and Schwartz FW. Sentinel practices in evaluating longer periods of care: quality of life and drug therapy of terminally ill persons in Lower Saxony (Germany). J Epidemiol Community Health 98;52 Suppl 1:56S-60S. [Rec#: 945]
INCLUDE: considered for evidence table

Butler JV, Pooviah PK, Cunningham D, and Hasan M. Improving decision-making and documentation relating to do not attempt resuscitation orders. Resuscitation 2003;57(2):139-44. [Rec#: 40494]
INCLUDE: considered for evidence table

Butow PN, Maclean M, Dunn SM, Tattersall, M. H, and Boyer, M. J. The dynamics of change: cancer patients' preferences for information, involvement and support. Annals of Oncology 97;8(9):857-63. [Rec#: 50312]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Cameron JI, Franche RL, Cheung AM, and Stewart DE. Lifestyle interference and emotional distress in family caregivers of advanced cancer patients. Cancer 2002;94(2):521-7. [Rec#: 985]
INCLUDE: considered for evidence table

Campbell ML and Guzman JA. Impact of a proactive approach to improve end-of-life care in a medical ICU. Chest 2003;123(1):266-71. Comment in: Chest. 2003 Jan;123(1):16-8. PMID: 12527596. [Rec#: 999]
INCLUDE: considered for evidence table

Campbell ML, Bizek KS, and Thill M. Patient responses during rapid terminal weaning from mechanical ventilation: a prospective study. Crit Care Med 99;27(1):73-7. Comment in: Crit Care Med. 1999 Jan;27(1):9-10. PMID: 9934874. [Rec#: 997]
INCLUDE: considered for evidence table

Carlson RW, Devich L, and Frank RR. Development of a comprehensive supportive care team for the hopelessly ill on a university hospital medical service. JAMA 88;259(3):378-83. [Rec#: 50321]
INCLUDE: considered for evidence table

Carson MG, Fitch MI, and Vachon ML. Measuring patient outcomes in palliative care: a reliability and validity study of the Support Team Assessment Schedule. Palliat Med 2000;14(1):25-36. [Rec#: 10067]
INCLUDE: considered for evidence table

Casarett DJ, Crowley R, and Hirschman KB. Surveys to assess satisfaction with end-of-life care: does timing matter? J Pain Symptom Manage 2003;25(2):128-32. [Rec#: 10091]
INCLUDE: considered for evidence table

Catalan-Fernandez JG, Pons-Sureda O, Recober-Martinez A, Avella-Mestre A, Carbonero-Malberti JM, Benito-Oliver E, and Garau-Llinas I. Dying of cancer. The place of death and family circumstances. Med Care 91;29(9):841-52. [Rec#: 10122]
INCLUDE: considered for evidence table

Centeno-Cortes C and Nunez-Olarte JM. Questioning diagnosis disclosure in terminal cancer patients: a prospective study evaluating patients' responses. Palliat Med 94;8(1):39-44. [Rec#: 10138]
INCLUDE: considered for evidence table

Chambers CV, Diamond JJ, Perkel RL, and Lasch LA. Relationship of advance directives to hospital charges in a Medicare population. Arch Intern Med 94;154(5):541-7. Comment in: Arch Intern Med. 1995 Jan 23;155(2):217-8. PMID: 7811135. Comment in: Arch Intern Med. 1995 Jan 9;155(1):116-7. PMID: 7802514. [Rec#: 1144]
INCLUDE: considered for evidence table

Chang VT, Thaler HT, Polyak TA, Kornblith AB, Lepore JM, and Portenoy RK. Quality of life and survival: the role of multidimensional symptom assessment. Cancer 98;83(1):173-9. [Rec#: 50333]
INCLUDE: considered for evidence table

Chelluri L, Im KA, Belle SH, Schulz R, Rotondi AJ, Donahoe MP, Sirio CA, Mendelsohn AB, and Pinsky MR. Long-term mortality and quality of life after prolonged mechanical ventilation. Crit Care Med 2004;32(1):61-9. Comment in: Crit Care Med. 2004 Jan;32(1):291-3. PMID: 14707597. [Rec#: 40593]
INCLUDE: considered for evidence table

Chen H, Haley WE, Robinson BE, and Schonwetter RS. Decisions for hospice care in patients with advanced cancer. J Am Geriatr Soc 2003;51(6):789-97. [Rec#: 1199]
INCLUDE: considered for evidence table

Chen JH, Bierhals AJ, Prigerson HG, Kasl SV, Mazure CM, and Jacobs S. Gender differences in the effects of bereavement-related psychological distress in health outcomes. Psychol Med 99;29(2):367-80. [Rec#: 1201]
INCLUDE: considered for evidence table

Chentsova-Dutton Y, Shuchter S, Hutchin S, Strause L, Burns K, and Zisook S. The psychological and physical health of hospice caregivers. Ann Clin Psychiatry 2000;12(1):19-27. [Rec#: 1211]
INCLUDE: considered for evidence table

Chin MH and Goldman L. Correlates of early hospital readmission or death in patients with congestive heart failure. Am J Cardiol 97;79(12):1640-4. [Rec#: 40615]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Chochinov HM, Hack T, Hassard T, Kristjanson LJ, McClement S, and Harlos M. Dignity in the terminally ill: a cross-sectional, cohort study. Lancet 2002;360(9350):2026-30. Comment in: Lancet. 2002 Dec 21-28;360(9350):1997-8. PMID: 12504390. Comment in: Lancet. 2003 Mar 1;361(9359):783. PMID: 12620760. [Rec#: 1252]
INCLUDE: considered for evidence table

Chochinov HM, Tataryn DJ, Wilson KG, Ennis M, and Lander S. Prognostic awareness and the terminally ill. Psychosomatics 2000;41(6):500-4. [Rec#: 1253]
INCLUDE: considered for evidence table

Chochinov HM, Wilson KG, Enns M, and Lander S. "Are you depressed?" Screening for depression in the terminally ill. Am J Psychiatry 97;154(5):674-6. Comment in: Am J Psychiatry. 1998 Jul;155(7):994-5. PMID: 9659876. [Rec#: 1254]
INCLUDE: considered for evidence table

Chochinov HM, Wilson KG, Enns M, and Lander S. Prevalence of depression in the terminally ill: effects of diagnostic criteria and symptom threshold judgments. Am J Psychiatry 94;151(4):537-40. [Rec#: 1256]
INCLUDE: considered for evidence table

Chow E, Andersson L, Wong R, Vachon M, Hruby G, Franssen E, Fung KW, Harth T, Pach B, Pope J, Connolly R, Schueller T, Stefaniuk K, Szumacher E, Hayter C, Finkelstein J, and Danjoux C. Patients with advanced cancer: a survey of the understanding of their illness and expectations from palliative radiotherapy for symptomatic metastases. Clin Oncol (R Coll Radiol) 2001;13(3):204-8. [Rec#: 1265]
INCLUDE: considered for evidence table

Chow E, Wong R, Connolly R, Hruby G, Franzcr, Franssen E, Fung KW, Vachon M, Andersson L, Pope J, Holden L, Szumacher E, Schueller T, Stefaniuk K, Finkelstein J, Hayter C, and Danjoux C. Prospective assessment of symptom palliation for patients attending a rapid response radiotherapy program. Feasibility of telephone follow-up. J Pain Symptom Manage 2001;22(2):649-56. [Rec#: 1268]
INCLUDE: considered for evidence table

Christakis NA and Iwashyna TJ. The health impact of health care on families: a matched cohort study of hospice use by decedents and mortality outcomes in surviving, widowed spouses. Soc Sci Med 2003;57(3):465-75. [Rec#: 1279]
INCLUDE: considered for evidence table

Christakis NA. Timing of referral of terminally ill patients to an outpatient hospice. J Gen Intern Med 94;9(6):314-20. [Rec#: 1276]
INCLUDE: considered for evidence table

Clark C and Heidenreich T. Spiritual care for the critically ill. Am J Crit Care 95;4(1):77-81. [Rec#: 40640]
INCLUDE: considered for evidence table

Clifford CA, Jolley DJ, and Giles GG. Where people die in Victoria. Med J Aust 91;155(7):446-51, 456. [Rec#: 1342]
INCLUDE: considered for evidence table

Clipp EC and George LK. Patients with cancer and their spouse caregivers. Perceptions of the illness experience. Cancer 92;69(4):1074-9. [Rec#: 50862]
INCLUDE: considered for evidence table

Cohen LM, Germain MJ, Poppel DM, Woods AL, Pekow PS, and Kjellstrand CM. Dying well after discontinuing the life-support treatment of dialysis. Arch Intern Med 2000;160(16):2513-8. [Rec#: 1372]
INCLUDE: considered for evidence table

Cohen LM, McCue JD, Germain M, and Woods A. Denying the dying. Advance directives and dialysis discontinuation. Psychosomatics 97;38(1):27-34. [Rec#: 1373]
INCLUDE: considered for evidence table

Cohen LM, McCue JD, Germain M, and Kjellstrand CM. Dialysis discontinuation. A 'good' death? Archives of Internal Medicine 95;155(1):42-7. [Rec#: 50344]
INCLUDE: considered for evidence table

Cohen SR, Mount BM, Tomas JJ, and Mount LF. Existential well-being is an important determinant of quality of life. Evidence from the McGill Quality of Life Questionnaire. Cancer 96;77(3):576-86. [Rec#: 50346]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Cohen-Mansfield J, Libin A, and Lipson S. Differences in presenting advance directives in the chart, in the minimum data set, and through the staff's perceptions. Gerontologist 2003;43(3):302-8. [Rec#: 1376]
INCLUDE: considered for evidence table

Colantonio A, Kositsky AJ, Cohen C, and Vernich L. What support do caregivers of elderly want? Results from the Canadian Study of Health and Aging. Can J Public Health 2001;92(5):376-9. [Rec#: 31020]
INCLUDE: considered for evidence table

Coll PP, Duffy JD, Micholovich E, and Cohen S. Best practices in hospital end-of-life care. Conn Med 2002;66(11):671-5. [Rec#: 10330]
INCLUDE: considered for evidence table

Collins C, Stommel M, Wang S, and Given CW. Caregiving transitions: changes in depression among family caregivers of relatives with dementia. Nurs Res 94;43(4):220-5. [Rec#: 1405]
INCLUDE: considered for evidence table

Colon M and Lyke J. Comparison of hospice use and demographics among European Americans, African Americans, and Latinos. Am J Hosp Palliat Care 2003;20(3):182-90. [Rec#: 1410]
INCLUDE: considered for evidence table

Conill C, Verger E, Henriquez I, Saiz N, Espier M, Lugo F, and Garrigos A. Symptom prevalence in the last week of life. J Pain Symptom Manage 97;14(6):328-31. [Rec#: 1422]
INCLUDE: considered for evidence table

Connolly M and O'Neill J. Teaching a research-based approach to the management of breathlessness in patients with lung cancer. Eur J Cancer Care (Engl) 99;8(1):30-6. [Rec#: 31056]
INCLUDE: considered for evidence table

Cools HJ, Berkhout AM, and De Bock GH. Subcutaneous morphine infusion by syringe driver for terminally ill patients. Age Ageing 96;25(3):206-8. [Rec#: 10391]
INCLUDE: considered for evidence table

Corner J, Halliday D, Haviland J, Douglas HR, Bath P, Clark D, Normand C, Beech N, Hughes P, Marples R, Seymour J, Skilbeck J, and Webb T. Exploring nursing outcomes for patients with advanced cancer following intervention by Macmillan specialist palliative care nurses. J Adv Nurs 2003;41(6):561-74. [Rec#: 1463]
INCLUDE: considered for evidence table

Correa-Velez I, Clavarino A, Barnett AG, and Eastwood H. Use of complementary and alternative medicine and quality of life: changes at the end of life. Palliat Med 2003;17(8):695-703. [Rec#: 1467]
INCLUDE: considered for evidence table

Costantini M, Balzi D, Garronec E, Orlandini C, Parodi S, Vercelli M, and Bruzzi P. Geographical variations of place of death among Italian communities suggest an inappropriate hospital use in the terminal phase of cancer disease. Public Health 2000;114(1):15-20. [Rec#: 1476]
INCLUDE: considered for evidence table

Costantini M, Higginson IJ, Boni L, Orengo MA, Garrone E, Henriquet F, and Bruzzi P. Effect of a palliative home care team on hospital admissions among patients with advanced cancer. Palliat Med 2003;17(4):315-21. [Rec#: 1478]
INCLUDE: considered for evidence table

Costantini M, Toscani F, Gallucci M, Brunelli C, Miccinesi G, Tamburini M, Paci E, Di Giulio P, Peruselli C, Higginson I, and Addington-Hall J. Terminal cancer patients and timing of referral to palliative care: a multicenter prospective cohort study. Italian Cooperative Research Group on Palliative Medicine. J Pain Symptom Manage 99;18(4):243-52. [Rec#: 1479]
INCLUDE: considered for evidence table

Covinsky KE, Fuller JD, Yaffe K, Johnston CB, Hamel MB, Lynn J, Teno JM, and Phillips RS. Communication and decision-making in seriously ill patients: findings of the SUPPORT project. The Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. J Am Geriatr Soc 2000;48(5 Suppl):S187-93. [Rec#: 1501]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Covinsky KE, Goldman L, Cook EF, Oye R, Desbiens N, Reding D, Fulkerson W, Connors AF Jr, Lynn J, and Phillips RS. The impact of serious illness on patients' families. SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. JAMA 94;272(23):1839-44. [Rec#: 40727]
INCLUDE: considered for evidence table

Covinsky KE, Kahana E, Chin MH, Palmer RM, Fortinsky RH, and Landefeld CS. Depressive symptoms and 3-year mortality in older hospitalized medical patients. Annals of Internal Medicine 99;130(7):563-9. [Rec#: 50359]
INCLUDE: considered for evidence table

Cowley AJ and Skene AM. Treatment of severe heart failure: quantity or quality of life? A trial of enoximone. Enoximone Investigators. Br Heart J 94;72(3):226-30. Comment in: Br Heart J. 1995 Jun;73(6):584. PMID: 7626364. [Rec#: 31102]
INCLUDE: considered for evidence table

Cullinane CA, Borneman T, Smith DD, Chu DZ, Ferrell BR, and Wagman LD. The surgical treatment of cancer: a comparison of resource utilization following procedures performed with a curative and palliative intent. Cancer 2003;98(10):2266-73. [Rec#: 1572]
INCLUDE: considered for evidence table

Curtis JR, Patrick DL, Caldwell ES, and Collier AC. The attitudes of patients with advanced AIDS toward use of the medical futility rationale in decisions to forego mechanical ventilation. Arch Intern Med 2000;160(11):1597-601. Comment in: Arch Intern Med. 2000 Jun 12;160(11):1565-6. PMID: 10847247. Comment in: Arch Intern Med. 2001 Jan 8;161(1):128-30. PMID: 11146716. [Rec#: 1586]
INCLUDE: considered for evidence table

Curtis JR, Patrick DL, Caldwell ES, and Collier AC. Why don't patients and physicians talk about end-of-life care? Barriers to communication for patients with acquired immunodeficiency syndrome and their primary care clinicians. Arch Intern Med 2000;160(11):1690-6. Comment in: Arch Intern Med. 2000 Jun 12;160(11):1565-6. PMID: 10847247. [Rec#: 1587]
INCLUDE: considered for evidence table

Curtisa JR, Patrick DL, Caldwell E, Greenlee H, and Collier AC. The quality of patient-doctor communication about end-of-life care: a study of patients with advanced AIDS and their primary care clinicians. AIDS 99;13(9):1123-31. [Rec#: 1585]
INCLUDE: considered for evidence table

Dales RE, O'Connor A, Hebert P, Sullivan K, McKim D, and Llewellyn-Thomas H. Intubation and mechanical ventilation for COPD: development of an instrument to elicit patient preferences. Chest 99;116(3):792-800. [Rec#: 1620]
INCLUDE: considered for evidence table

Danis M, Southerland LI, Garrett JM, Smith JL, Hielema F, Pickard CG, Egner DM, and Patrick DL. A prospective study of advance directives for life-sustaining care. N Engl J Med 91;324(13):882-8. Comment in: N Engl J Med. 1991 Oct 24;325(17):1254-6. PMID: 1922218. [Rec#: 40771]
INCLUDE: considered for evidence table

Davies HT, Crombie IK, Brown JH, and Martin C. Diminishing returns or appropriate treatment strategy?--an analysis of short-term outcomes after pain clinic treatment. Pain 97;70(2-3):203-8. [Rec#: 1671]
INCLUDE: considered for evidence table

Davison BJ, Gleave ME, Goldenberg SL, Degner LF, Hoffart D, and Berkowitz J. Assessing information and decision preferences of men with prostate cancer and their partners. Cancer Nursing 2002;25(1):42-9. [Rec#: 50368]
INCLUDE: considered for evidence table

Dawson NJ. Need satisfaction in terminal care settings. Soc Sci Med 91;32(1):83-7. [Rec#: 10637]
INCLUDE: considered for evidence table

De Conno F, Boffi R, Brunelli C, and Panzeri C. Age-related differences in patients admitted to a palliative home care service. Tumori 2002;88(2):117-22. [Rec#: 1702]
INCLUDE: considered for evidence table

Appendix M. Included Studies

De Conno F, Caraceni A, Groff L, Brunelli C, Donati I, Tamburini M, and Ventafridda V. Effect of home care on the place of death of advanced cancer patients. Eur J Cancer 96;32A(7):1142-7. [Rec#: 1703]
INCLUDE: considered for evidence table

De Conno F, Saita L, Ripamonti C, and Ventafridda V. Subcutaneous octreotide in the treatment of pain in advanced cancer patients. J Pain Symptom Manage 94;9(1):34-8. [Rec#: 31225]
INCLUDE: considered for evidence table

De Jonge KE, Sulmasy DP, Gold KG, Epstein A, Harper MG, Eisenberg JM, and Schulman KA. The timing of do-not-resuscitate orders and hospital costs. J Gen Intern Med 99;14(3):190-2. [Rec#: 1712]
INCLUDE: considered for evidence table

de Vos R, Koster RW, and de Haan RJ. Impact of survival probability, life expectancy, quality of life and patient preferences on do-not-attempt-resuscitation orders in a hospital. Resuscitation Committee. Resuscitation 98;39(1-2):15-21. [Rec#: 40833]
INCLUDE: considered for evidence table

de Wit R, van Dam F, Abu-Saad HH, Loonstra S, Zandbelt L, van Buuren A, van der Heijden K, and Leenhouts G. Empirical comparison of commonly used measures to evaluate pain treatment in cancer patients with chronic pain. J Clin Oncol 99;17(4):1280. [Rec#: 1733]
INCLUDE: considered for evidence table

de Wit R, van Dam F, Vielvoye-Kerkmeer A, Mattern C, and Abu-Saad HH. The treatment of chronic cancer pain in a cancer hospital in The Netherlands. J Pain Symptom Manage 99;17(5):333-50. [Rec#: 10666]
INCLUDE: considered for evidence table

DeCourtney CA, Jones K, Merriman MP, Heavener N, and Branch PK. Establishing a culturally sensitive palliative care program in rural Alaska Native American communities. J Palliat Med 2003;6(3):501-10. [Rec#: 10682]
INCLUDE: considered for evidence table

Degenholtz HB, Arnold RA, Meisel A, and Lave JR. Persistence of racial disparities in advance care plan documents among nursing home residents. J Am Geriatr Soc 2002;50(2):378-81. [Rec#: 40841]
INCLUDE: considered for evidence table

Delbridge TR, Fosnocht DE, Garrison HG, and Auble TE. Field termination of unsuccessful out-of-hospital cardiac arrest resuscitation: acceptance by family members. Ann Emerg Med 96;27(5):649-54. [Rec#: 1757]
INCLUDE: considered for evidence table

Dendaas N, Pellino TA, Roberts KF, and Cleary J. Responding to SUPPORT: an academic medical center examines its end-of-life care practices. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. J Pain Symptom Manage 2001;21(2):121-8. [Rec#: 1777]
INCLUDE: considered for evidence table

Desbiens NA, Mueller-Rizner N, Connors AF Jr, Wenger NS, and Lynn J. The symptom burden of seriously ill hospitalized patients. SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcome and Risks of Treatment. J Pain Symptom Manage 99;17(4):248-55. [Rec#: 40862]
INCLUDE: considered for evidence table

Desbiens NA, Mueller-Rizner N, Hamel MB, and Connors AF Jr. Preference for comfort care does not affect the pain experience of seriously ill patients. The SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. J Pain Symptom Manage 98;16(5):281-9. [Rec#: 1787]
INCLUDE: considered for evidence table

Detmar SB, Muller MJ, Schornagel JH, Wever LD, and Aaronson NK. Health-related quality-of-life assessments and patient-physician communication: a randomized controlled trial. JAMA 2002;288(23):3027-34. Comment in: JAMA. 2003 Apr 2;289(13):1636; author reply 1636-7. PMID: 12672729. Erratum in: JAMA. 2003 Feb 26;289(8):987. [Rec#: 1794]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Detmar SB, Muller MJ, Schornagel JH, Wever LD, and Aaronson NK. Role of health-related quality of life in palliative chemotherapy treatment decisions. J Clin Oncol 2002;20(4):1056-62. [Rec#: 1795]
INCLUDE: considered for evidence table

Detmar SB, Muller MJ, Wever LD, Schornagel JH, and Aaronson NK. The patient-physician relationship. Patient-physician communication during outpatient palliative treatment visits: an observational study. JAMA 2001;285(10):1351-7. [Rec#: 1796]
INCLUDE: considered for evidence table

Devery K, Lennie I, and Cooney N. Health outcomes for people who use palliative care services. J Palliat Care 99;15(2):5-12. [Rec#: 10719]
INCLUDE: considered for evidence table

Devulder J, Ghys L, Dhondt W, and Rolly G. Spinal analgesia in terminal care: risk versus benefit. J Pain Symptom Manage 94;9(2):75-81. [Rec#: 10730]
INCLUDE: considered for evidence table

Dexter PR, Wolinsky FD, Gramelspacher GP, Zhou XH, Eckert GJ, Waisburd M, and Tierney WM. Effectiveness of computer-generated reminders for increasing discussions about advance directives and completion of advance directive forms. A randomized, controlled trial. Ann Intern Med 98;128(2):102-10. [Rec#: 50914]
INCLUDE: considered for evidence table

Ditto PH, Danks JH, Smucker WD, Bookwala J, Coppola KM, Dresser R, Fagerlin A, Gready RM, Houts RM, Lockhart LK, and Zyzanski S. Advance directives as acts of communication: a randomized controlled trial. Arch Intern Med 2001;161(3):421-30. [Rec#: 1852]
INCLUDE: considered for evidence table

Dobratz MC. Analysis of variables that impact psychological adaptation in home hospice patients. Hosp J 95;10(1):75-88. [Rec#: 1865]
INCLUDE: considered for evidence table

Dobratz MC. Patterns of advanced cancer pain in home hospice patients. Cancer Nurs 2001;24(4):294-9. [Rec#: 1870]
INCLUDE: considered for evidence table

Donaldson N, Carter H, and Green R. Quality of information on hospice referral. Br J Gen Pract 2000;50(452):219-20. [Rec#: 10793]
INCLUDE: considered for evidence table

Donnelly S and Walsh D. The symptoms of advanced cancer. Semin Oncol 95;22(2 Suppl 3):67-72. [Rec#: 1896]
INCLUDE: considered for evidence table

Donovan KA, Greene PG, Shuster JL, Partridge EE, and Tucker DC. Treatment preferences in recurrent ovarian cancer. Gynecol Oncol 2002;86(2):200-11. [Rec#: 1899]
INCLUDE: considered for evidence table

Doorenbos AZ and Nies MA. The use of advance directives in a population of Asian Indian Hindus. J Transcult Nurs 2003;14(1):17-24. [Rec#: 1903]
INCLUDE: considered for evidence table

Dowdy MD, Robertson C, and Bander JA. A study of proactive ethics consultation for critically and terminally ill patients with extended lengths of stay. Critical Care Medicine 98;26(2):252-9. [Rec#: 50384]
INCLUDE: considered for evidence table

Dowsett SM, Saul JL, Butow PN, Dunn SM, Boyer MJ, Findlow R, and Dunsmore J. Communication styles in the cancer consultation: preferences for a patient-centred approach. Psycho-Oncology 2000;9(2):147-56. [Rec#: 50385]
INCLUDE: considered for evidence table

Doyle CJ. Effect of staff turnover and the social environment on depressive symptoms in nursing home residents. Int Psychogeriatr 95;7(1):51-61. [Rec#: 40917]
INCLUDE: considered for evidence table

Draper B, MacCuspie-Moore C, and Brodaty H. Suicidal ideation and the 'wish to die' in dementia patients: the role of depression. Age Ageing 98;27(4):503-7. [Rec#: 1934]
INCLUDE: considered for evidence table

Dudgeon DJ and Lertzman M. Dyspnea in the advanced cancer patient. J Pain Symptom Manage 98;16(4):212-9. Comment in: J Pain Symptom Manage. 1999 Nov;18(5):313-5. PMID: 10584452. [Rec#: 40930]
INCLUDE: considered for evidence table

Dudgeon DJ, Raubertas RF, Doerner K, O'Connor T, Tobin M, and Rosenthal SN. When does palliative care begin? A needs assessment of cancer patients with recurrent disease. J Palliat Care 95;11(1):5-9. Comment in: J Palliat Care. 1995 Autumn;11(3):53. PMID: 7472797. Erratum in: J Palliat Care. 1995 Autumn;11(3):53. PMID: 7472797. [Rec#: 1955]
INCLUDE: considered for evidence table

Duggleby W and Alden C. Implementation and evaluation of a quality improvement process to improve pain management in a hospice setting. Am J Hosp Palliat Care 98;15(4):209-16. [Rec#: 10867]
INCLUDE: considered for evidence table

Dull SM, Graves JR, Larsen MP, and Cummins RO. Expected death and unwanted resuscitation in the prehospital setting. Ann Emerg Med 94;23(5):997-1002. [Rec#: 40939]
INCLUDE: considered for evidence table

Dumot JA, Burval DJ, Sprung J, Waters JH, Mraovic B, Karafa MT, Mascha EJ, and Bourke DL. Outcome of adult cardiopulmonary resuscitations at a tertiary referral center including results of "limited" resuscitations. Arch Intern Med 2001;161(14):1751-8. [Rec#: 40940]
INCLUDE: considered for evidence table

Duthie E, Mark D, Tresch D, Kartes S, Neahring J, and Aufderheide T. Utilization of cardiopulmonary resuscitation in nursing homes in one community: rates and nursing home characteristics. J Am Geriatr Soc 93;41(4):384-8. Comment in: J Am Geriatr Soc. 1993 Nov;41(11):1275-6. PMID: 8227908. [Rec#: 40956]
INCLUDE: considered for evidence table

DuVal G, Clarridge B, Gensler G, and Danis M. A national survey of U.S. internists' experiences with ethical dilemmas and ethics consultation. J Gen Intern Med 2004;19(3):251-8. [Rec#: 1989]
INCLUDE: considered for evidence table

Ehman JW, Ott BB, Short TH, Ciampa RC, and Hansen-Flaschen J. Do patients want physicians to inquire about their spiritual or religious beliefs if they become gravely ill? Archives of Internal Medicine 99;159(15):1803-6. [Rec#: 50396]
INCLUDE: considered for evidence table

Eisdorfer C, Czaja, SJ, Loewenstein DA, Rubert MP, Arguelles S, Mitrani VB, and Szapocznik J. The effect of a family therapy and technology-based intervention on caregiver depression. Gerontologist 2003;43(4):521-31. [Rec#: 50917]
INCLUDE: considered for evidence table

Eldridge AD, Severance-Lossin L, Nicholas PK, and Leuner JD. Prevalence and characteristics of pain in persons with terminal-stage AIDS. J Adv Nurs 94;20(2):260-8. [Rec#: 2059]
INCLUDE: considered for evidence table

Eleazer GP, Hornung CA, Egbert CB, Egbert JR, Eng C, Hedgepeth J, McCann R, Strothers H 3rd, Sapir M, Wei M, and Wilson M. The relationship between ethnicity and advance directives in a frail older population. J Am Geriatr Soc 96;44(8):938-43. [Rec#: 40988]
INCLUDE: considered for evidence table

Elfrink EJ, van der Rijt CC, van Boxtel RJ, Elswijk-de Vries P, van Zuijlen L, and Stoter G. Problem solving by telephone in palliative care: use of a predetermined assessment tool within a program of home care technology. J Palliat Care 2002;18(2):105-10. [Rec#: 2061]
INCLUDE: considered for evidence table

Elmstahl S, Ingvad B, and Annerstedt L. Family caregiving in dementia: prediction of caregiver burden 12 months after relocation to group-living care. Int Psychogeriatr 98;10(2):127-46. [Rec#: 50026]
INCLUDE: considered for evidence table

Emanuel EJ, Fairclough DL, Slutsman J, Alpert H, Baldwin D, and Emanuel LL. Assistance from family members, friends, paid care givers, and volunteers in the care of terminally ill patients. N Engl J Med 99;341(13):956-63. [Rec#: 2092]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Emanuel EJ, Weinberg DS, Gonin R, Hummel LR, and Emanuel LL. How well is the Patient Self-Determination Act working?: an early assessment. Am J Med 93;95(6):619-28. [Rec#: 2093]
INCLUDE: considered for evidence table

Emanuel EJ, Fairclough DL, Slutsman J, and Emanuel LL. Understanding economic and other burdens of terminal illness: the experience of patients and their caregivers. Annals of Internal Medicine 2000;132(6):451-9. [Rec#: 50398]
INCLUDE: considered for evidence table

Emanuel LL, Barry MJ, Stoeckle JD, Ettelson LM, and Emanuel EJ. Advance directives for medical care—a case for greater use. N Engl J Med 91;324(13):889-95. [Rec#: 50864]
INCLUDE: considered for evidence table

Escalante CP, Martin CG, Elting LS, Cantor SB, Harle TS, Price KJ, Kish SK, Manzullo EF, and Rubenstein EB. Dyspnea in cancer patients. Etiology, resource utilization, and survival—implications in a managed care world. Cancer 96;78(6):1314-9. [Rec#: 2131]
INCLUDE: considered for evidence table

Evangelista LS, Dracup K, Doering L, Westlake C, Fonarow GC, and Hamilton M. Emotional well-being of heart failure patients and their caregivers. J Card Fail 2002;8(5):300-5. [Rec#: 31516]
INCLUDE: considered for evidence table

Evers MM, Meier DE, and Morrison RS. Assessing differences in care needs and service utilization in geriatric palliative care patients. J Pain Symptom Manage 2002;23(5):424-32. [Rec#: 2157]
INCLUDE: considered for evidence table

Evers MM, Purohit D, Perl D, Khan K, and Marin DB. Palliative and aggressive end-of-life care for patients with dementia. Psychiatr Serv 2002;53(5):609-13. [Rec#: 2158]
INCLUDE: considered for evidence table

Faber-Langendoen K and Bartels DM. Process of forgoing life-sustaining treatment in a university hospital: an empirical study. Crit Care Med 92;20(5):570-7. [Rec#: 2163]
INCLUDE: considered for evidence table

Faber-Langendoen, K. A multi-institutional study of care given to patients dying in hospitals. Ethical and practice implications. Archives of Internal Medicine 96;156(18):2130-6. [Rec#: 50403]
INCLUDE: considered for evidence table

Fainsinger R, Miller MJ, Bruera E, Hanson J, and Maceachern T. Symptom control during the last week of life on a palliative care unit. J Palliat Care 91;7(1):5-11. [Rec#: 2171]
INCLUDE: considered for evidence table

Fainsinger RL, De Moissac D, Mancini I, and Oneschuk D. Sedation for delirium and other symptoms in terminally ill patients in Edmonton. J Palliat Care 2000;16(2):5-10. [Rec#: 2172]
INCLUDE: considered for evidence table

Fainsinger RL, Louie K, Belzile M, Bruera E, and Hanson J. Decreased opioid doses used on a palliative care unit. J Palliat Care 96;12(4):6-9. [Rec#: 2176]
INCLUDE: considered for evidence table

Fainsinger RL, MacEachern T, Hanson J, and Bruera E. The use of urinary catheters in terminally ill cancer patients. J Pain Symptom Manage 92;7(6):333-8. [Rec#: 2177]
INCLUDE: considered for evidence table

Fainsinger RL, MacEachern T, Miller MJ, Bruera E, Spachynski K, Kuehn N, and Hanson J. The use of hypodermoclysis for rehydration in terminally ill cancer patients. J Pain Symptom Manage 94;9(5):298-302. [Rec#: 2178]
INCLUDE: considered for evidence table

Fainsinger RL, Nunez-Olarte JM, and Demoissac DM. The cultural differences in perceived value of disclosure and cognition: Spain and Canada. J Palliat Care 2003;19(1):43-8. Comment in: J Palliat Care. 2003 Spring;19(1):7-8. PMID: 12710109. [Rec#: 2179]
INCLUDE: considered for evidence table

Fainsinger RL, Spachynski K, Hanson J, and Bruera E. Symptom control in terminally ill patients with malignant bowel obstruction (MBO). J Pain Symptom Manage 94;9(1):12-8. [Rec#: 2181]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Fakhoury W, McCarthy M, and Addington-Hall J. Determinants of informal caregivers' satisfaction with services for dying cancer patients. Soc Sci Med 96;42(5):721-31. [Rec#: 2190]
INCLUDE: considered for evidence table

Fakhoury WK, McCarthy M, and Addington-Hall J. Carers' health status: is it associated with their evaluation of the quality of palliative care? Scand J Soc Med 97;25(4):296-301. [Rec#: 2192]
INCLUDE: considered for evidence table

Fakhoury WK, McCarthy M, and Addington-Hall J. The effects of the clinical characteristics of dying cancer patients on informal caregivers' satisfaction with palliative care. Palliat Med 97;11(2):107-15. [Rec#: 2193]
INCLUDE: considered for evidence table

Farran CJ, Miller BH, Kaufman JE, Donner E, and Fogg L. Finding meaning through caregiving: development of an instrument for family caregivers of persons with Alzheimer's disease. J Clin Psychol 99;55(9):1107-25. [Rec#: 31551]
INCLUDE: considered for evidence table

Fernsler JI, Klemm P, and Miller MA. Spiritual well-being and demands of illness in people with colorectal cancer. Cancer Nurs 99;22(2):134-40; quiz 141-2. [Rec#: 2253]
INCLUDE: considered for evidence table

Ferrell BA, Ferrell BR, and Rivera L. Pain in cognitively impaired nursing home patients. J Pain Symptom Manage 95;10(8):591-8. [Rec#: 41059]
INCLUDE: considered for evidence table

Ferry P, Hancock R, Newton C, Taylor J, and Horton K. Perspectives of relatives and health care workers on care of terminally ill patients in the UK. Am J Hosp Palliat Care 2002;19(2):121-8. [Rec#: 11176]
INCLUDE: considered for evidence table

Feuz A and Rapin CH. An observational study of the role of pain control and food adaptation of elderly patients with terminal cancer. J Am Diet Assoc 94;94(7):767-70. [Rec#: 2279]
INCLUDE: considered for evidence table

Field D and McGaughey J. An evaluation of palliative care services for cancer patients in the Southern Health and Social Services Board of Northern Ireland. Palliat Med 98;12(2):83-97. [Rec#: 2286]
INCLUDE: considered for evidence table

Fine RL and Mayo TW. Resolution of futility by due process: early experience with the Texas Advance Directives Act. Ann Intern Med 2003;138(9):743-6. Comment in: Ann Intern Med. 2004 Mar 2;140(5):402-3; author reply 404. PMID: 14996685. Comment in: Ann Intern Med. 2004 Mar 2;140(5):403-4; author reply 404. PMID: 14996687. Comment in: Ann Intern Med. 2004 Mar 2;140(5):403; author reply 404. PMID. [Rec#: 2304]
INCLUDE: considered for evidence table

Finlay IG and Jones OL. Hypnotherapy in palliative care. J R Soc Med 96;89(9):493-6. [Rec#: 2311]
INCLUDE: considered for evidence table

Finn JW, Walsh TD, MacDonald N, Bruera E, Krebs LU, and Shepard KV. Placebo-blinded study of morphine sulfate sustained-release tablets and immediate-release morphine sulfate solution in outpatients with chronic pain due to advanced cancer. Journal of Clinical Oncology 93;11(5):967-72. [Rec#: 50428]
INCLUDE: considered for evidence table

Fins JJ, Miller FG, Acres CA, Bacchetta MD, Huzzard LL, and Rapkin BD. End-of-life decision-making in the hospital: current practice and future prospects. J Pain Symptom Manage 99;17(1):6-15. [Rec#: 2314]
INCLUDE: considered for evidence table

Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, and Pinder EL. The implications of regional variations in Medicare spending. Part 2: health outcomes and satisfaction with care. Ann Intern Med 2003;138(4):288-98. Comment in: Ann Intern Med. 2003 Feb 18;138(4):347-8. PMID: 12585834. Comment in: Ann Intern Med. 2003 Feb 18;138(4):348-9. PMID: 12585835. Comment in: Ann Intern Med. 2003 Feb 18;138(4):350-1. PMID: 12585836. Comment in: Ann Intern Med. 2004 Jan . [Rec#: 2326]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Fitzpatrick MO, Goyal K, and Barlow P. Treatment limiting decisions in a neurosurgical unit. Scott Med J 99;44(2):47-8. [Rec#: 11252]
INCLUDE: considered for evidence table

Fogel BS and Mor V. Depressed mood and care preferences in patients with AIDS. Gen Hosp Psychiatry 93;15(4):203-7. [Rec#: 2360]
INCLUDE: considered for evidence table

Fortner BV, Demarco G, Irving G, Ashley J, Keppler G, Chavez J, and Munk J. Description and predictors of direct and indirect costs of pain reported by cancer patients. J Pain Symptom Manage 2003;25(1):9-18. [Rec#: 11303]
INCLUDE: considered for evidence table

Fountain A. Visual hallucinations: a prevalence study among hospice inpatients. Palliat Med 2001;15(1):19-25. [Rec#: 2388]
INCLUDE: considered for evidence table

Fowell A, Finlay I, Johnstone R, and Minto L. An integrated care pathway for the last two days of life: Wales-wide benchmarking in palliative care. Int J Palliat Nurs 2002;8(12):566-73. Comment in: Int J Palliat Nurs. 2003 Jan;9(1):39. PMID: 12560796. [Rec#: 11314]
INCLUDE: considered for evidence table

Fried TR and Bradley EH. What matters to seriously ill older persons making end-of-life treatment decisions?: A qualitative study. J Palliat Med 2003;6(2):237-44. [Rec#: 2435]
INCLUDE: considered for evidence table

Fried TR and Mor V. Frailty and hospitalization of long-term stay nursing home residents. J Am Geriatr Soc 97;45(3):265-9. Comment in: J Am Geriatr Soc. 1997 Mar;45(3):378-9. PMID: 9063288. [Rec#: 41137]
INCLUDE: considered for evidence table

Fried TR, Bradley EH, and O'Leary J. Prognosis communication in serious illness: perceptions of older patients, caregivers, and clinicians. J Am Geriatr Soc 2003;51(10):1398-403. [Rec#: 2436]
INCLUDE: considered for evidence table

Fried TR, Bradley EH, and Towle VR. Assessment of patient preferences: integrating treatments and outcomes. J Gerontol B Psychol Sci Soc Sci 2002;57(6):S348-54. [Rec#: 2437]
INCLUDE: considered for evidence table

Fried TR, Bradley EH, and Towle VR. Valuing the outcomes of treatment: do patients and their caregivers agree? Arch Intern Med 2003;163(17):2073-8. [Rec#: 2438]
INCLUDE: considered for evidence table

Fried TR, Bradley EH, Towle VR, and Allore H. Understanding the treatment preferences of seriously ill patients. N Engl J Med 2002;346(14):1061-6. Comment in: N Engl J Med. 2002 Apr 4;346(14):1087-9. PMID: 11932479. Comment in: N Engl J Med. 2002 Aug 15;347(7):533-5; author reply 533-5. PMID: 12181411. Comment in: N Engl J Med. 2002 Aug 15;347(7):533-5; author reply 533-5. PMID: 12182184. Co. [Rec#: 2439]
INCLUDE: considered for evidence table

Fried TR, Pollack DM, Drickamer MA, and Tinetti ME. Who dies at home? Determinants of site of death for community-based long-term care patients. J Am Geriatr Soc 99;47(1):25-9. [Rec#: 2441]
INCLUDE: considered for evidence table

Fried TR, van Doorn C, O'Leary JR, Tinetti ME, and Drickamer MA. Older person's preferences for home vs hospital care in the treatment of acute illness. Arch Intern Med 2000;160(10):1501-6. [Rec#: 41139]
INCLUDE: considered for evidence table

Fried TR, Gillick MR, and Lipsitz L. A. Short-term functional outcomes of long-term care residents with pneumonia treated with and without hospital transfer. Journal of the American Geriatrics Society 97;45(3):302-6. [Rec#: 50437]
INCLUDE: considered for evidence table

Friedman MM and Griffin JA. Relationship of physical symptoms and physical functioning to depression in patients with heart failure. Heart Lung 2001;30(2):98-104. [Rec#: 31732]
INCLUDE: considered for evidence table

Friedrichsen MJ, Strang PM, and Carlsson ME. Cancer patients' perceptions of their participation and own resources after receiving information about discontinuation of active tumour treatment. Acta Oncol 2000;39(8):919-25. [Rec#: 2451]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Fry PS. A factor analytic investigation of home-bound elderly individuals' concerns about death and dying, and their coping responses. J Clin Psychol 90;46(6):737-48. [Rec#: 2469]
INCLUDE: considered for evidence table

Fudin J, Smith HS, Toledo-Binette CS, Kenney E, Yu AB, and Boutin R. Use of continuous ambulatory infusions of concentrated subcutaneous (s.q.) hydromorphone versus intravenous (i.v.) morphine: cost implications for palliative care. Am J Hosp Palliat Care 2000;17(5):347-53. Comment in: Am J Hosp Palliat Care. 2000 Sep-Oct;17(5):292-3. PMID: 11886046. [Rec#: 11399]
INCLUDE: considered for evidence table

Fulop G, Strain JJ, and Stettin G. Congestive heart failure and depression in older adults: clinical course and health services use 6 months after hospitalization. Psychosomatics 2003;44(5):367-73. [Rec#: 31762]
INCLUDE: considered for evidence table

Fulton C. Patients with metastatic breast cancer: their physical and psychological rehabilitation needs. Int J Rehabil Res 99;22(4):291-301. [Rec#: 31763]
INCLUDE: considered for evidence table

Fulton CL. The physical and psychological symptoms experienced by patients with metastatic breast cancer before death. Eur J Cancer Care (Engl) 97;6(4):262-6. [Rec#: 2489]
INCLUDE: considered for evidence table

Furuta M, Hayakawa K, Saito Y, Nakayama Y, Katano S, Sakurai H, Takahashi T, Ohno T, Mitsuhashi N, and Niibe H. Clinical implication of symptoms in patients with non-small cell lung cancer treated with definitive radiation therapy. Lung Cancer 95;13(3):275-83. [Rec#: 31769]
INCLUDE: considered for evidence table

Gagnon P, Allard P, Masse B, and DeSerres M. Delirium in terminal cancer: a prospective study using daily screening, early diagnosis, and continuous monitoring. J Pain Symptom Manage 2000;19(6):412-26. [Rec#: 2512]
INCLUDE: considered for evidence table

Gagnon, B, Bielech, M, Watanabe, S, Walker, P, Hanson, J, and Bruera, E. The use of intermittent subcutaneous injections of oxycodone for opioid rotation in patients with cancer pain. Supportive Care in Cancer 99;7(4):265-70. [Rec#: 50440]
INCLUDE: considered for evidence table

Gallicchio L, Siddiqi N, Langenberg P, and Baumgarten M. Gender differences in burden and depression among informal caregivers of demented elders in the community. Int J Geriatr Psychiatry 2002;17(2):154-63. [Rec#: 31785]
INCLUDE: considered for evidence table

Gallo WT, Baker MJ, and Bradley EH. Factors associated with home versus institutional death among cancer patients in Connecticut. J Am Geriatr Soc 2001;49(6):771-7. Comment in: J Am Geriatr Soc. 2001 Jun;49(6):831-2. PMID: 11454126. [Rec#: 2527]
INCLUDE: considered for evidence table

Ganzini L, Lee MA, Heintz RT, and Bloom JD. Do-not-resuscitate orders for depressed psychiatric inpatients. Hosp Community Psychiatry 92;43(9):915-9. [Rec#: 2543]
INCLUDE: considered for evidence table

Ganzini L, Goy ER, Miller LL, Harvath TA, Jackson A, and Delorit MA. Nurses' experiences with hospice patients who refuse food and fluids to hasten death. New England Journal of Medicine 2003;349(4):359-65. [Rec#: 50444]
INCLUDE: considered for evidence table

Ganzini L, Johnston WS, and Hoffman WF. Correlates of suffering in amyotrophic lateral sclerosis. Neurology 99;52(7):1434-40. [Rec#: 50445]
INCLUDE: considered for evidence table

Gattellari M, Voigt KJ, Butow PN, and Tattersall MH. When the treatment goal is not cure: are cancer patients equipped to make informed decisions? J Clin Oncol 2002;20(2):503-13. [Rec#: 2565]
INCLUDE: considered for evidence table

Gaudette LA, Shi F, Lipskie T, Allard P, Fainsinger RL, Maxwell D, and Harlos M. Developing palliative care surveillance in Canada: results of a pilot study. J Palliat Care 2002;18(4):262-9. [Rec#: 2567]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Ghusn HF, Teasdale TA, and Jordan D. Continuity of do-not-resuscitate orders between hospital and nursing home settings. J Am Geriatr Soc 97;45(4):465-9. [Rec#: 41196]
INCLUDE: considered for evidence table

Gift AG, Stommel M, Jablonski A, and Given W. A cluster of symptoms over time in patients with lung cancer. Nurs Res 2003;52(6):393-400. [Rec#: 2633]
INCLUDE: considered for evidence table

Giladi N, Treves TA, Paleacu D, Shabtai H, Orlov Y, Kandinov B, Simon ES, and Korczyn AD. Risk factors for dementia, depression and psychosis in long-standing Parkinson's disease. J Neural Transm 2000;107(1):59-71. [Rec#: 31892]
INCLUDE: considered for evidence table

Gill P, Kaur JS, Rummans T, Novotny PJ, and Sloan JA. The hospice patient's primary caregiver. What is their quality of life? J Psychosom Res 2003;55(5):445-51. [Rec#: 2643]
INCLUDE: considered for evidence table

Giorgi F, Cellerino R, Gramazio A, Tummarello D, Menichetti ET, Giordani P, Antognoli S, Carle F, and Piga A. Assessing quality of life in patients with cancer: a comparison of a visual-analogue and a categorical model. Am J Clin Oncol 96;19(4):394-9. [Rec#: 31911]
INCLUDE: considered for evidence table

Gitlin LN, Belle SH, Burgio LD, Czaja SJ, Mahoney D, Gallagher-Thompson D, Burns R, Hauck WW, Zhang S, Schulz R, Ory MG, and REACH Investigators. Effect of multicomponent interventions on caregiver burden and depression: the REACH multisite initiative at 6-month follow-up. Psychol Aging 2003;18(3):361-74. [Rec#: 31919]
INCLUDE: considered for evidence table

Gitlin LN, Winter L, Corcoran M, Dennis MP, Schinfeld S, and Hauck WW. Effects of the home environmental skill-building program on the caregiver-care recipient dyad: 6-month outcomes from the Philadelphia REACH Initiative. Gerontologist 2003;43(4):532-46. [Rec#: 50918]
INCLUDE: considered for evidence table

Given B, Given C, Azzouz F, and Stommel M. Physical functioning of elderly cancer patients prior to diagnosis and following initial treatment. Nurs Res 2001;50(4):222-32. [Rec#: 31922]
INCLUDE: considered for evidence table

Gochman DS and Bonham GS. The social structure of the hospice decision. Hosp J 90;6(1):15-36. [Rec#: 11560]
INCLUDE: considered for evidence table

Goldberg LR, Piette JD, Walsh MN, Frank TA, Jaski BE, Smith AL, Rodriguez R, Mancini DM, Hopton LA, Orav EJ, Loh E, and WHARF Investigators. Randomized trial of a daily electronic home monitoring system in patients with advanced heart failure: the Weight Monitoring in Heart Failure (WHARF) trial. Am Heart J 2003;146(4):705-12. [Rec#: 31947]
INCLUDE: considered for evidence table

Goldberg RJ and Goldberg J. Risperidone for dementia-related disturbed behavior in nursing home residents: a clinical experience. Int Psychogeriatr 97;9(1):65-8. [Rec#: 31950]
INCLUDE: considered for evidence table

Goldstone I, Kuhl D, Johnson A, Le R, and McLeod A. Patterns of care in advanced HIV disease in a tertiary treatment centre. AIDS Care 95;7 Suppl 1:S47-56. [Rec#: 2714]
INCLUDE: considered for evidence table

Golin CE, Wenger NS, Liu H, Dawson NV, Teno JM, Desbiens NA, Lynn J, Oye RK, and Phillips RS. A prospective study of patient-physician communication about resuscitation. J Am Geriatr Soc 2000;48(5 Suppl):S52-60. [Rec#: 2716]
INCLUDE: considered for evidence table

Goncalves JF, Alvarenga M, and Silva A. The last forty-eight hours of life in a Portuguese palliative care unit: does it differ from elsewhere? J Palliat Med 2003;6(6):895-900. [Rec#: 2720]
INCLUDE: considered for evidence table

Gonzalez-Salvador MT, Arango C, Lyketsos CG, and Barba AC. The stress and psychological morbidity of the Alzheimer patient caregiver. Int J Geriatr Psychiatry 99;14(9):701-10. [Rec#: 31971]
INCLUDE: considered for evidence table

Goodkin K, Feaster DJ, Asthana D, Blaney NT, Kumar M, Baldewicz T, Tuttle RS, Maher KJ, Baum MK, Shapshak P, and Fletcher MA. A bereavement support group intervention is longitudinally associated with salutary effects on the CD4 cell count and number of physician visits. Clin Diagn Lab Immunol 98;5(3):382-91. [Rec#: 2725]
INCLUDE: considered for evidence table

Goodlin SJ, Winzelberg GS, Teno JM, Whedon M, and Lynn J. Death in the hospital. Arch Intern Med 98;158(14):1570-2. [Rec#: 2729]
INCLUDE: considered for evidence table

Goodlin SJ, Zhong Z, Lynn J, Teno JM, Fago JP, Desbiens N, Connors AF Jr, Wenger NS, and Phillips RS. Factors associated with use of cardiopulmonary resuscitation in seriously ill hospitalized adults. JAMA 99;282(24):2333-9. [Rec#: 2730]
INCLUDE: considered for evidence table

Goodwin DM, Higginson IJ, Myers K, Douglas HR, and Normand CE. Effectiveness of palliative day care in improving pain, symptom control, and quality of life. J Pain Symptom Manage 2003;25(3):202-12. Comment in: J Pain Symptom Manage. 2003 Oct;26(4):886-7; author reply 888-9. PMID: 14527753. [Rec#: 2735]
INCLUDE: considered for evidence table

Goodwin DM, Higginson IJ, Myers K, Douglas HR, and Normand CE. What is palliative day care? A patient perspective of five UK services. Support Care Cancer 2002;10(7):556-62. [Rec#: 2736]
INCLUDE: considered for evidence table

Goodyer LI, Miskelly F, and Milligan P. Does encouraging good compliance improve patients' clinical condition in heart failure? Br J Clin Pract 95;49(4):173-6. Comment in: Br J Clin Pract. 1995 Jul-Aug;49(4):171. PMID: 7547152. [Rec#: 31978]
INCLUDE: considered for evidence table

Gordon NP and Shade SB. Advance directives are more likely among seniors asked about end-of-life care preferences. Arch Intern Med 99;159(7):701-4. [Rec#: 2742]
INCLUDE: considered for evidence table

Gore JM, Brophy CJ, and Greenstone MA. How well do we care for patients with end stage chronic obstructive pulmonary disease (COPD)? A comparison of palliative care and quality of life in COPD and lung cancer. Thorax 2000;55(12):1000-6. Comment in: Thorax. 2000 Dec;55(12):979-81. PMID: 11083879. [Rec#: 2745]
INCLUDE: considered for evidence table

Gorski LA and Johnson K. A disease management program for heart failure: collaboration between a home care agency and a care management organization. Lippincotts Case Manag 2003;8(6):265-73. [Rec#: 31989]
INCLUDE: considered for evidence table

Gottlieb SS, Fisher ML, Freudenberger R, Robinson S, Zietowski G, Alves L, Krichten C, Vaitkevicus P, and McCarter R. Effects of exercise training on peak performance and quality of life in congestive heart failure patients. J Card Fail 99;5(3):188-94. [Rec#: 31996]
INCLUDE: considered for evidence table

Grande GE, McKerral A, and Todd CJ. Which cancer patients are referred to Hospital at Home for palliative care? Palliat Med 2002;16(2):115-23. [Rec#: 2777]
INCLUDE: considered for evidence table

Grande GE, Todd CJ, Barclay SI, and Farquhar MC. A randomized controlled trial of a hospital at home service for the terminally ill. Palliat Med 2000;14(5):375-85. [Rec#: 11650]
INCLUDE: considered for evidence table

Grande GE, Todd CJ, Barclay SI, and Farquhar MC. Does hospital at home for palliative care facilitate death at home? Randomised controlled trial. BMJ 99;319(7223):1472-5. Comment in: BMJ. 1999 Dec 4;319(7223):1447-8. PMID: 10582909. [Rec#: 11649]
INCLUDE: considered for evidence table

Grant I, Adler KA, Patterson TL, Dimsdale JE, Ziegler MG, and Irwin MR. Health consequences of Alzheimer's caregiving transitions: effects of placement and bereavement. Psychosom Med 2002;64(3):477-86. [Rec#: 2781]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Grant M, Golant M, Rivera L, Dean G, and Benjamin H. Developing a community program on cancer pain and fatigue. Cancer Practice 2000;8(4):187-94. [Rec#: 50465]
INCLUDE: considered for evidence table

Green CP, Porter CB, Bresnahan DR, and Spertus JA. Development and evaluation of the Kansas City Cardiomyopathy Questionnaire: a new health status measure for heart failure. J Am Coll Cardiol 2000;35(5):1245-55. [Rec#: 32030]
INCLUDE: considered for evidence table

Green CR and Wheeler JR. Physician variability in the management of acute postoperative and cancer pain: a quantitative analysis of the Michigan experience. Pain Med 2003;4(1):8-20. Comment in: Pain Med. 2003 Mar;4(1):1-3. PMID: 12873272. [Rec#: 11670]
INCLUDE: considered for evidence table

Greenberg SM, Tennis MK, Brown LB, Gomez-Isla T, Hayden DL, Schoenfeld DA, Walsh KL, Corwin C, Daffner KR, Friedman P, Meadows ME, Sperling RA, and Growdon JH. Donepezil therapy in clinical practice: a randomized crossover study. Arch Neurol 2000;57(1):94-9. Comment in: Arch Neurol. 2000 Sep;57(9):1380. PMID: 10987910. [Rec#: 32039]
INCLUDE: considered for evidence table

Griffith CH 3rd, Wilson JF, Emmett KR, Ramsbottom-Lucier M, and Rich EC. Knowledge and experience with Alzheimer's disease. Relationship to resuscitation preference. Arch Fam Med 95;4(9):780-4. [Rec#: 32062]
INCLUDE: considered for evidence table

Groff L, Zecca E, De Conno F, Brunelli C, Boffi R, Panzeri C, Cazzaniga M, and Ripamonti C. The role of disodium pamidronate in the management of bone pain due to malignancy. Palliat Med 2001;15(4):297-307. [Rec#: 2834]
INCLUDE: considered for evidence table

Grond S, Zech D, Lynch J, Diefenbach C, Schug SA, and Lehmann KA. Validation of World Health Organization guidelines for pain relief in head and neck cancer. A prospective study. Ann Otol Rhinol Laryngol 93;102(5):342-8. [Rec#: 2837]
INCLUDE: considered for evidence table

Gross MD. What do patients express as their preferences in advance directives? Arch Intern Med 98;158(4):363-5. Comment in: Arch Intern Med. 1998 Feb 23;158(4):321-4. PMID: 9487228. [Rec#: 11731]
INCLUDE: considered for evidence table

Guo H, Fine PG, Mendoza TR, and Cleeland CS. A preliminary study of the utility of the brief hospice inventory. J Pain Symptom Manage 2001;22(2):637-48. [Rec#: 2870]
INCLUDE: considered for evidence table

Guthrie B, Nelson M, and Gazzard B. Are people with HIV in London able to die where they plan? AIDS Care 96;8(6):709-13. [Rec#: 11769]
INCLUDE: considered for evidence table

Gutzmann H, Kuhl KP, Hadler D, and Rapp MA. Safety and efficacy of idebenone versus tacrine in patients with Alzheimer's disease: results of a randomized, double-blind, parallel-group multicenter study. Pharmacopsychiatry 2002;35(1):12-8. [Rec#: 32110]
INCLUDE: considered for evidence table

Guyatt GH, Berman LB, Townsend M, Pugsley SO, and Chambers LW. A measure of quality of life for clinical trials in chronic lung disease. Thorax 87;42(10):773-8. [Rec#: 50867]
INCLUDE: considered for evidence table

Haggmark C, Bachner M, and Theorell T. A follow-up of psychological state in relatives of cancer patients one year after the patient's death. Effects of an activation programme. Acta Oncol 91;30(6):677-84. [Rec#: 2911]
INCLUDE: considered for evidence table

Hakim RB, Teno JM, Harrell FE Jr, Knaus WA, Wenger N, Phillips RS, Layde P, Califf R, Connors AF Jr, and Lynn J. Factors associated with do-not-resuscitate orders: patients' preferences, prognoses, and physicians' judgments. SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. Ann Intern Med 96;125(4):284-93. [Rec#: 41337]
INCLUDE: considered for evidence table

Haley WE, LaMonde LA, Han B, Burton AM, and Schonwetter R. Predictors of depression and life satisfaction among spousal caregivers in hospice: application of a stress process model. J Palliat Med 2003;6(2):215-24. [Rec#: 2937]
INCLUDE: considered for evidence table

Hall P, Schroder C, and Weaver L. The last 48 hours of life in long-term care: a focused chart audit. J Am Geriatr Soc 2002;50(3):501-6. [Rec#: 2945]
INCLUDE: considered for evidence table

Hall S, Gallagher RM, Gracely E, Knowlton C, and Wescules D. The terminal cancer patient: effects of age, gender, and primary tumor site on opioid dose. Pain Med 2003;4(2):125-34. [Rec#: 2947]
INCLUDE: considered for evidence table

Hamel MB, Lynn J, Teno JM, Covinsky KE, Wu AW, Galanos A, Desbiens NA, and Phillips RS. Age-related differences in care preferences, treatment decisions, and clinical outcomes of seriously ill hospitalized adults: lessons from SUPPORT. J Am Geriatr Soc 2000;48(5 Suppl):S176-82. [Rec#: 2954]
INCLUDE: considered for evidence table

Hamel MB, Phillips RS, Davis RB, Teno J, Connors AF, Desbiens N, Lynn J, Dawson NV, Fulkerson W, and Tsevat J. Outcomes and cost-effectiveness of ventilator support and aggressive care for patients with acute respiratory failure due to pneumonia or acute respiratory distress syndrome. American Journal of Medicine 2000;109(8):614-20. [Rec#: 50474]
INCLUDE: considered for evidence table

Hamel MB, Phillips R S, Teno JM, Lynn J, Galanos AN, Davis RB, Connors AF Jr, Oye RK, Desbiens N, Reding DJ, and Goldman L. Seriously ill hospitalized adults: do we spend less on older patients? Support Investigators. Study to Understand Prognoses and Preference for Outcomes and Risks of Treatments. Journal of the American Geriatrics Society 96;44(9):1043-8. [Rec#: 50476]
INCLUDE: considered for evidence table

Hammes BJ and Rooney BL. Death and end-of-life planning in one midwestern community. Arch Intern Med 98;158(4):383-90. [Rec#: 2961]
INCLUDE: considered for evidence table

Hanks GW, Robbins M, Sharp D, Forbes K, Done K, Peters TJ, Morgan H, Sykes J, Baxter K, Corfe F, and Bidgood C. The imPaCT study: a randomised controlled trial to evaluate a hospital palliative care team. Br J Cancer 2002;87(7):733-9. [Rec#: 2978]
INCLUDE: considered for evidence table

Hanrahan P and Luchins DJ. Access to hospice programs in end-stage dementia: a national survey of hospice programs. J Am Geriatr Soc 95;43(1):56-9. Comment in: J Am Geriatr Soc. 1995 Oct;43(10):1174. PMID: 7560715. Comment in: J Am Geriatr Soc. 1995 Oct;43(10):1174-6. PMID: 7560716. [Rec#: 11846]
INCLUDE: considered for evidence table

Hanson LC and Rodgman E. The use of living wills at the end of life. A national study. Arch Intern Med 96;156(9):1018-22. [Rec#: 11861]
INCLUDE: considered for evidence table

Hanson LC, Earp JA, Garrett J, Menon M, and Danis M. Community physicians who provide terminal care. Arch Intern Med 99;159(10):1133-8. [Rec#: 2992]
INCLUDE: considered for evidence table

Harding R, Higginson IJ, and Donaldson N. The relationship between patient characteristics and carer psychological status in home palliative cancer care. Support Care Cancer 2003;11(10):638-43. [Rec#: 3008]
INCLUDE: considered for evidence table

Hardy JR, Rees E, Ling J, Burman R, Feuer D, Broadley K, and Stone P. A prospective survey of the use of dexamethasone on a palliative care unit. Palliat Med 2001;15(1):3-8. Comment in: Palliat Med. 2001 Nov;15(6):521. PMID: 12403513. [Rec#: 11877]
INCLUDE: considered for evidence table

Hardy JR, Rees EA, Gwilliam B, Ling J, Broadley K, and A'Hern R. A phase II study to establish the efficacy and toxicity of sodium valproate in patients with cancer-related neuropathic pain. J Pain Symptom Manage 2001;21(3):204-9. Comment in: J Pain Symptom Manage. 2002 Jan;23(1):1. PMID: 11779658. [Rec#: 3015]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Harrell LE, Callaway R, Morere D, and Falgout J. The effect of long-term physostigmine administration in Alzheimer's disease. Neurology 90;40(9):1350-4. [Rec#: 32179]
INCLUDE: considered for evidence table

Harrison J, Haddad P, and Maguire P. The impact of cancer on key relatives: a comparison of relative and patient concerns. European Journal of Cancer 95;31A(11):1736-40. [Rec#: 50483]
INCLUDE: considered for evidence table

Haydar ZR, Lowe AJ, Kahveci KL, Weatherford W, and Finucane T. Differences in end-of-life preferences between congestive heart failure and dementia in a medical house calls program. J Am Geriatr Soc 2004;52(5):736-40. [Rec#: 50854]
INCLUDE: considered for evidence table

Hays JC, Kasl SV, and Jacobs SC. The course of psychological distress following threatened and actual conjugal bereavement. Psychol Med 94;24(4):917-27. [Rec#: 3084]
INCLUDE: considered for evidence table

Heap MJ, Munglani R, Klinck JR, and Males AG. Elderly patients' preferences concerning life-support treatment. Anaesthesia 93;48(12):1027-33. Comment in: Anaesthesia. 1993 Dec;48(12):1025-6. PMID: 8285319. [Rec#: 3095]
INCLUDE: considered for evidence table

Heaven CM and Maguire P. The relationship between patients' concerns and psychological distress in a hospice setting. Psychooncology 98;7(6):502-7. [Rec#: 3097]
INCLUDE: considered for evidence table

Heedman PA and Strang P. Pain and pain alleviation in hospital-based home care: demographic, biological and treatment factors. Support Care Cancer 2003;11(1):35-40. [Rec#: 3103]
INCLUDE: considered for evidence table

Heeren O, Menon AS, Raskin A, and Ruskin P. Religion and end of life treatment preferences among geriatric patients. Int J Geriatr Psychiatry 2001;16(2):203-8. [Rec#: 3105]
INCLUDE: considered for evidence table

Heffner JE and Barbieri C. End-of-life care preferences of patients enrolled in cardiovascular rehabilitation programs. Chest 2000;117(5):1474-81. Comment in: Chest. 2000 May;117(5):1228-31. PMID: 10807804. [Rec#: 3107]
INCLUDE: considered for evidence table

Heffner JE and Barbieri C. Involvement of cardiovascular rehabilitation programs in advance directive education. Arch Intern Med 96;156(15):1746-51. [Rec#: 11989]
INCLUDE: considered for evidence table

Heffner JE, Barbieri C, and Casey K. Procedure-specific do-not-resuscitate orders. Effect on communication of treatment limitations. Arch Intern Med 96;156(7):793-7. [Rec#: 41447]
INCLUDE: considered for evidence table

Heffner JE, Barbieri C, Fracica P, and Brown LK. Communicating do-not-resuscitate orders with a computer-based system. Arch Intern Med 98;158(10):1090-5. [Rec#: 41448]
INCLUDE: considered for evidence table

Heffner JE, Fahy B, Hilling L, and Barbieri C. Attitudes regarding advance directives among patients in pulmonary rehabilitation. Am J Respir Crit Care Med 96;154(6 Pt 1):1735-40. [Rec#: 3108]
INCLUDE: considered for evidence table

Heffner JE, Fahy B, Hilling L, and Barbieri C. Outcomes of advance directive education of pulmonary rehabilitation patients. Am J Respir Crit Care Med 97;155(3):1055-9. [Rec#: 3109]
INCLUDE: considered for evidence table

Heidenreich PA, Ruggerio CM, and Massie BM. Effect of a home monitoring system on hospitalization and resource use for patients with heart failure. Am Heart J 99;138(4 Pt 1):633-40. Comment in: Am Heart J. 1999 Oct;138(4 Pt 1):599-601. PMID: 10502199. [Rec#: 32251]
INCLUDE: considered for evidence table

Helft PR, Hlubocky F, Wen M, and Daugherty CK. Associations among awareness of prognosis, hopefulness, and coping in patients with advanced cancer participating in phase I clinical trials. Support Care Cancer 2003;11(10):644-51. [Rec#: 41456]
INCLUDE: considered for evidence table

Heller RF, Fisher JD, D'Este CA, Lim LL, Dobson AJ, and Porter R. Death and readmission in the year after hospital admission with cardiovascular disease: the Hunter Area Heart and Stroke Register. Med J Aust 2000;172(6):261-5. [Rec#: 50036]
INCLUDE: considered for evidence table

Herndon JE 2nd, Fleishman S, Kosty MP, and Green MR. A longitudinal study of quality of life in advanced non-small cell lung cancer: Cancer and Leukemia Group B (CALGB) 8931. Control Clin Trials 97;18(4):286-300. Comment in: Control Clin Trials. 1997 Aug;18(4):306-10. PMID: 9257069. Comment in: Control Clin Trials. 1997 Aug;18(4):311-7. PMID: 9257070. [Rec#: 32303]
INCLUDE: considered for evidence table

Hesse KA. Terminal care of the very old. Changes in the way we die. Arch Intern Med 95;155(14):1513-8. [Rec#: 3180]
INCLUDE: considered for evidence table

Heyland DK, Konopad E, Noseworthy TW, Johnston R, and Gafni A. Is it 'worthwhile' to continue treating patients with a prolonged stay (>14 days) in the ICU? An economic evaluation. Chest 98;114(1):192-8. [Rec#: 41500]
INCLUDE: considered for evidence table

Heyland DK, Lavery JV, Tranmer JE, Shortt SE, and Taylor SJ. Dying in Canada: is it an institutionalized, technologically supported experience? J Palliat Care 2000;16 Suppl:S10-6. [Rec#: 12051]
INCLUDE: considered for evidence table

Heyland DK, Rocker GM, O'Callaghan CJ, Dodek PM, and Cook DJ. Dying in the ICU: perspectives of family members. Chest 2003;124(1):392-7. Comment in: Chest. 2003 Jul;124(1):11-2. PMID: 12853494. [Rec#: 3184]
INCLUDE: considered for evidence table

Heyland DK, Tranmer J, O'Callaghan CJ, and Gafni A. The seriously ill hospitalized patient: preferred role in end-of-life decision making? J Crit Care 2003;18(1):3-10. [Rec#: 3186]
INCLUDE: considered for evidence table

Hickok JT, Morrow GR, McDonald S, and Bellg AJ. Frequency and correlates of fatigue in lung cancer patients receiving radiation therapy: implications for management. J Pain Symptom Manage 96;11(6):370-7. [Rec#: 32326]
INCLUDE: considered for evidence table

Higginson I and Priest P. Predictors of family anxiety in the weeks before bereavement. Soc Sci Med 96;43(11):1621-5. [Rec#: 3198]
INCLUDE: considered for evidence table

Higginson I, Priest P, and McCarthy M. Are bereaved family members a valid proxy for a patient's assessment of dying? Soc Sci Med 94;38(4):553-7. [Rec#: 12063]
INCLUDE: considered for evidence table

Higginson IJ and Costantini M. Communication in end-of-life cancer care: a comparison of team assessments in three European countries. J Clin Oncol 2002;20(17):3674-82. [Rec#: 3202]
INCLUDE: considered for evidence table

Hill KM, Amir Z, Muers MF, Connolly CK, and Round CE. Do newly diagnosed lung cancer patients feel their concerns are being met? Eur J Cancer Care (Engl) 2003;12(1):35-45. [Rec#: 32331]
INCLUDE: considered for evidence table

Hilliard RE. The effects of music therapy on the quality and length of life of people diagnosed with terminal cancer. J Music Ther 2003;40(2):113-37. [Rec#: 3213]
INCLUDE: considered for evidence table

Hinds C. Personal and contextual factors predicting patients' reported quality of life: exploring congruency with Betty Neuman's assumptions. J Adv Nurs 90;15(4):456-62. [Rec#: 32340]
INCLUDE: considered for evidence table

Hines SC, Glover JJ, Babrow AS, Holley JL, Badzek LA, and Moss AH. Improving advance care planning by accommodating family preferences. J Palliat Med 2001;4(4):481-9. Comment in: J Palliat Med. 2001 Winter;4(4):453-6. PMID: 11798476. [Rec#: 3226]
INCLUDE: considered for evidence table

Hines SC, Glover JJ, Holley JL, Babrow AS, Badzek LA, and Moss AH. Dialysis patients' preferences for family-based advance care planning. Ann Intern Med 99;130(10):825-8. [Rec#: 3227]
INCLUDE: considered for evidence table

Hinkka H, Kosunen E, Kellokumpu-Lehtinen P, and Lammi UK. Assessment of pain control in cancer patients during the last week of life: comparison of health centre wards and a hospice. Support Care Cancer 2001;9(6):428-34. Comment in: Support Care Cancer. 2001 Sep;9(6):401-2. PMID: 11585265. [Rec#: 3228]
INCLUDE: considered for evidence table

Hinton J. Can home care maintain an acceptable quality of life for patients with terminal cancer and their relatives? Palliat Med 94;8(3):183-96. [Rec#: 3236]
INCLUDE: considered for evidence table

Hinton J. How reliable are relatives' retrospective reports of terminal illness? Patients and relatives' accounts compared. Soc Sci Med 96;43(8):1229-36. [Rec#: 3237]
INCLUDE: considered for evidence table

Hinton J. Services given and help perceived during home care for terminal cancer. Palliat Med 96;10(2):125-34. [Rec#: 3239]
INCLUDE: considered for evidence table

Hinton J. Which patients with terminal cancer are admitted from home care? Palliat Med 94;8(3):197-210. [Rec#: 3240]
INCLUDE: considered for evidence table

Hodgson C, Higginson I, McDonnell M, and Butters E. Family anxiety in advanced cancer: a multicentre prospective study in Ireland. Br J Cancer 97;76(9):1211-4. [Rec#: 3261]
INCLUDE: considered for evidence table

Hodgson H. Does reflexology impact on cancer patients' quality of life? Nurs Stand 2000;14(31):33-8. [Rec#: 3262]
INCLUDE: considered for evidence table

Holden CM. Anorexia in the terminally ill cancer patient: the emotional impact on the patient and the family. Hosp J 91;7(3):73-84. [Rec#: 3285]
INCLUDE: considered for evidence table

Holden JD. Auditing palliative care in one general practice over eight years. Scand J Prim Health Care 96;14(3):136-41. [Rec#: 12149]
INCLUDE: considered for evidence table

Hollen PJ, Gralla RJ, Kris MG, Eberly SW, and Cox C. Normative data and trends in quality of life from the Lung Cancer Symptom Scale (LCSS). Support Care Cancer 99;7(3):140-8. [Rec#: 32384]
INCLUDE: considered for evidence table

Holley JL, Hines SC, Glover JJ, Babrow AS, Badzek LA, and Moss AH. Failure of advance care planning to elicit patients' preferences for withdrawal from dialysis. Am J Kidney Dis 99;33(4):688-93. [Rec#: 3291]
INCLUDE: considered for evidence table

Holtzman J, Chen Q, and Kane R. The effect of HMO status on the outcomes of home-care after hospitalization in a Medicare population. J Am Geriatr Soc 98;46(5):629-34. [Rec#: 50040]
INCLUDE: considered for evidence table

Holtzman J, Pheley AM, and Lurie N. Changes in orders limiting care and the use of less aggressive care in a nursing home population. J Am Geriatr Soc 94;42(3):275-9. [Rec#: 41540]
INCLUDE: considered for evidence table

Holzapfel L, Demingeon G, Piralla B, Biot L, and Nallet B. A four-step protocol for limitation of treatment in terminal care. An observational study in 475 intensive care unit patients. Intensive Care Med 2002;28(9):1309-15. Comment in: Intensive Care Med. 2002 Sep;28(9):1197-9. PMID: 12400561. [Rec#: 3303]
INCLUDE: considered for evidence table

Holzheimer A, McMillan SC, and Weitzner M. Improving pain outcomes of hospice patients with cancer. Oncol Nurs Forum 99;26(9):1499-504. [Rec#: 3304]
INCLUDE: considered for evidence table

Homsi J, Walsh D, Nelson KA, LeGrand SB, Davis M, Khawam E, and Nouneh C. The impact of a palliative medicine consultation service in medical oncology. Support Care Cancer 2002;10(4):337-42. [Rec#: 3311]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Hopwood P and Stephens RJ. Depression in patients with lung cancer: prevalence and risk factors derived from quality-of-life data. J Clin Oncol 2000;18(4):893-903. [Rec#: 3332]
INCLUDE: considered for evidence table

Hopwood P, Harvey A, Davies J, Stephens RJ, Girling DJ, Gibson D, and Parmar MK. Survey of the Administration of quality of life (QL) questionnaires in three multicentre randomised trials in cancer. The Medical Research Council Lung Cancer Working Party the CHART Steering Committee. Eur J Cancer 98;34(1):49-57. [Rec#: 32409]
INCLUDE: considered for evidence table

Hopwood P, Howell A, and Maguire P. Psychiatric morbidity in patients with advanced cancer of the breast: prevalence measured by two self-rating questionnaires. British Journal of Cancer 91;64(2):349-52. [Rec#: 50499]
INCLUDE: considered for evidence table

Hopwood P, Howell A, and Maguire P. Screening for psychiatric morbidity in patients with advanced breast cancer: validation of two self-report questionnaires. British Journal of Cancer 91;64(2):353-6. [Rec#: 50500]
INCLUDE: considered for evidence table

Houts RM, Smucker WD, Jacobson JA, Ditto PH, and Danks JH. Predicting elderly outpatients' life-sustaining treatment preferences over time: the majority rules. Med Decis Making 2002;22(1):39-52. [Rec#: 3364]
INCLUDE: considered for evidence table

Hugel H, Ellershaw JE, Cook L, Skinner J, and Irvine C. The prevalence, key causes and management of insomnia in palliative care patients. J Pain Symptom Manage 2004;27(4):316-21. [Rec#: 3394]
INCLUDE: considered for evidence table

Hughes SL, Weaver FM, Giobbie-Hurder A, Manheim L, Henderson W, Kubal JD, Ulasevich A, Cummings J, and Department of Veterans Affairs Cooperative Study Group on Home-Based Primary Care. Effectiveness of team-managed home-based primary care: a randomized multicenter trial. JAMA 2000;284(22):2877-85. [Rec#: 3399]
INCLUDE: considered for evidence table

Hunt R, Fazekas B, Thorne D, and Brooksbank M. A comparison of subcutaneous morphine and fentanyl in hospice cancer patients. J Pain Symptom Manage 99;18(2):111-9. [Rec#: 3422]
INCLUDE: considered for evidence table

Hunt RW, Fazekas BS, Luke CG, and Roder DM. Where patients with cancer die in South Australia, 1990-1999: a population-based review. Med J Aust 2001;175(10):526-9. Comment in: Med J Aust. 2001 Nov 19;175(10):534-6. PMID: 11795543. [Rec#: 3426]
INCLUDE: considered for evidence table

Hutt E, Ecord M, Eilertsen TB, Frederickson E, and Kramer AM. Precipitants of emergency room visits and acute hospitalization in short-stay medicare nursing home residents. J Am Geriatr Soc 2002;50(2):223-9. [Rec#: 32478]
INCLUDE: considered for evidence table

Hwang SS, Chang VT, Fairclough DL, and Kasimis B. Development of a cancer pain prognostic scale. J Pain Symptom Manage 2002;24(4):366-78. [Rec#: 3448]
INCLUDE: considered for evidence table

Ibbotson T, Maguire P, Selby P, Priestman T, and Wallace L. Screening for anxiety and depression in cancer patients: the effects of disease and treatment. European Journal of Cancer 94;30A(1):37-40. [Rec#: 50502]
INCLUDE: considered for evidence table

Im K, Belle SH, Schulz R, Mendelsohn AB, Chelluri L, and QOL-MV Investigators. Prevalence and outcomes of caregiving after prolonged (> or =48 hours) mechanical ventilation in the ICU. Chest 2004;125(2):597-606. [Rec#: 41609]
INCLUDE: considered for evidence table

Incalzi RA, Gemma A, Marra C, Capparella O, Fuso L, and Carbonin P. Verbal memory impairment in COPD: its mechanisms and clinical relevance. Chest 97;112(6):1506-13. [Rec#: 32517]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Ingleton C, Morgan J, Hughes P, Noble B, Evans A, and Clark D. Carer satisfaction with end-of-life care in Powys, Wales: a cross-sectional survey. Health Soc Care Community 2004;12(1):43-52. [Rec#: 3473]
INCLUDE: considered for evidence table

Jaagosild P, Dawson NV, Thomas C, Wenger NS, Tsevat J, Knaus WA, Califf RM, Goldman L, Vidaillet H, and Connors AF Jr. Outcomes of acute exacerbation of severe congestive heart failure: quality of life, resource use, and survival. SUPPORT Investigators. The Study to Understand Prognosis and Preferences for Outcomes and Risks of Treatments. Arch Intern Med 98;158(10):1081-9. [Rec#: 32575]
INCLUDE: considered for evidence table

Jaarsma T, Dracup K, Walden J, and Stevenson LW. Sexual function in patients with advanced heart failure. Heart Lung 96;25(4):262-70. [Rec#: 32580]
INCLUDE: considered for evidence table

Jaarsma T, Halfens R, Huijer Abu-Saad H, Dracup K, Gorgels T, van Ree J, and Stappers J. Effects of education and support on self-care and resource utilization in patients with heart failure. Eur Heart J 99;20(9):673-82. Comment in: Eur Heart J. 1999 May;20(9):632-3. PMID: 10419340. [Rec#: 32582]
INCLUDE: considered for evidence table

Jaarsma T, Halfens R, Tan F, Abu-Saad HH, Dracup K, and Diederiks J. Self-care and quality of life in patients with advanced heart failure: the effect of a supportive educational intervention. Heart Lung 2000;29(5):319-30. [Rec#: 32583]
INCLUDE: considered for evidence table

Jaarsma T. Sexual problems in heart failure patients. Eur J Cardiovasc Nurs 2002;1(1):61-7. [Rec#: 32578]
INCLUDE: considered for evidence table

Jack B, Hillier V, Williams A, and Oldham J. Hospital based palliative care teams improve the insight of cancer patients into their disease. Palliat Med 2004;18(1):46-52. [Rec#: 3510]
INCLUDE: considered for evidence table

Jack B, Hillier V, Williams A, and Oldham J. Hospital based palliative care teams improve the symptoms of cancer patients. Palliat Med 2003;17(6):498-502. [Rec#: 3511]
INCLUDE: considered for evidence table

Jacobs LG, Bonuck K, and Burton W. Can "Palliative Care Reports" improve end-of-life care for hospitalized patients? J Pain Symptom Manage 2002;24(3):299-311. [Rec#: 3530]
INCLUDE: considered for evidence table

Jacobs S, Hansen F, Kasl S, Ostfeld A, Berkman L, and Kim K. Anxiety disorders during acute bereavement: risk and risk factors. J Clin Psychiatry 90;51(7):269-74. Comment in: J Clin Psychiatry. 1991 May;52(5):241. PMID: 2033034. [Rec#: 3534]
INCLUDE: considered for evidence table

Jacobson JA, Kasworm E, Battin MP, Francis LP, Green D, Botkin J, and Johnson S. Advance directives in Utah. Information from death certificates and informants. Arch Intern Med 96;156(16):1862-8. [Rec#: 41636]
INCLUDE: considered for evidence table

Jacoby A, Lecouturier J, Bradshaw C, Lovel T, and Eccles M. Feasibility of using postal questionnaires to examine carer satisfaction with palliative care: a methodological assessment. South Tyneside MAAG Palliative Care Study Group. Palliat Med 99;13(4):285-98. [Rec#: 12351]
INCLUDE: considered for evidence table

Janjan NA, Payne R, Gillis T, Podoloff D, Libshitz HI, Lenzi R, Theriault R, Martin C, and Yasko A. Presenting symptoms in patients referred to a multidisciplinary clinic for bone metastases. J Pain Symptom Manage 98;16(3):171-8. [Rec#: 3561]
INCLUDE: considered for evidence table

Jayes RL, Zimmerman JE, Wagner DP, and Knaus WA. Variations in the use of do-not-resuscitate orders in ICUS. Findings from a national study. Chest 96;110(5):1332-9. Comment in: Chest. 1996 Nov;110(5):1141-2. PMID: 8915210. [Rec#: 12388]
INCLUDE: considered for evidence table

Jenkins CA, Schulz M, Hanson J, and Bruera E. Demographic, symptom, and medication profiles of cancer patients seen by a palliative care consult team in a tertiary referral hospital. J Pain Symptom Manage 2000;19(3):174-84. [Rec#: 3578]
INCLUDE: considered for evidence table

Jenkins CA, Taube AW, Turner K, Hanson J, and Bruera E. Initial demographic, symptom, and medication profiles in patients admitted to continuing palliative care units. J Pain Symptom Manage 98;16(3):163-70. [Rec#: 3579]
INCLUDE: considered for evidence table

Jerant AF, Azari R, and Nesbitt TS. Reducing the cost of frequent hospital admissions for congestive heart failure: a randomized trial of a home telecare intervention. Med Care 2001;39(11):1234-45. [Rec#: 50047]
INCLUDE: considered for evidence table

Jerant AF, Azari RS, Nesbitt TS, and Meyers FJ. The TLC model of palliative care in the elderly: preliminary application in the assisted living setting. Ann Fam Med 2004;2(1):54-60. [Rec#: 3591]
INCLUDE: considered for evidence table

Jezewski MA and Finnell DS. The meaning of DNR status: oncology nurses' experiences with patients and families. Cancer Nurs 98;21(3):212-21. [Rec#: 3599]
INCLUDE: considered for evidence table

Jiang W, Alexander J, Christopher E, Kuchibhatla M, Gaulden LH, Cuffe MS, Blazing MA, Davenport C, Califf RM, Krishnan RR, and O'Connor CM. Relationship of depression to increased risk of mortality and rehospitalization in patients with congestive heart failure. Arch Intern Med 2001;161(15):1849-56. Comment in: Arch Intern Med. 2002 Feb 11;162(3):362. PMID: 11822938. Comment in: Arch Intern Med. 2002 Feb 11;162(3):362-4. PMID: 11822939. [Rec#: 50048]
INCLUDE: considered for evidence table

Johnson IS. Use of bisphosphonates for the treatment of metastatic bone pain. A survey of palliative physicians in the UK. Palliat Med 2001;15(2):141-7. [Rec#: 12440]
INCLUDE: considered for evidence table

Johnson MF, Lin M, Mangalik S, Murphy DJ, and Kramer AM. Patients' perceptions of physicians' recommendations for comfort care differ by patient age and gender. J Gen Intern Med 2000;15(4):248-55. [Rec#: 3624]
INCLUDE: considered for evidence table

Johnston G and Abraham C. The WHO objectives for palliative care: to what extent are we achieving them? Palliat Med 95;9(2):123-37. [Rec#: 12456]
INCLUDE: considered for evidence table

Johnston SC and Johnson SC. Advance directives: from the perspective of the patient and the physician. J R Soc Med 96;89(10):568-70. Erratum in: J R Soc Med 1997 Feb;90(2):120. Johnson SC[corrected to Johnston SC]. [Rec#: 12461]
INCLUDE: considered for evidence table

Johnstone PA, Polston GR, Niemtzow RC, and Martin PJ. Integration of acupuncture into the oncology clinic. Palliat Med 2002;16(3):235-9. [Rec#: 3635]
INCLUDE: considered for evidence table

Jordhoy MS, Fayers P, Loge JH, Ahlner-Elmqvist M, and Kaasa S. Quality of life in palliative cancer care: results from a cluster randomized trial. J Clin Oncol 2001;19(18):3884-94. Comment in: J Clin Oncol. 2002 Mar 15;20(6):1704-5. PMID: 11896122. [Rec#: 3660]
INCLUDE: considered for evidence table

Jordhoy MS, Saltvedt I, Fayers P, Loge JH, Ahlner-Elmqvist M, and Kaasa S. Which cancer patients die in nursing homes? Quality of life, medical and sociodemographic characteristics. Palliat Med 2003;17(5):433-44. [Rec#: 3664]
INCLUDE: considered for evidence table

Jorm AF, Henderson S, Scott R, Mackinnon AJ, Korten AE, and Christensen H. The disabled elderly living in the community: care received from family and formal services. Med J Aust 93;158(6):383-5, 388. [Rec#: 41697]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Junod Perron N, Morabia A, and de Torrente A. Quality of life of Do-Not-Resuscitate (DNR) patients: how good are physicians in assessing DNR patients' quality of life? Swiss Med Wkly 2002;132(39-40):562-5. [Rec#: 41704]
INCLUDE: considered for evidence table

Kaasa S, Malt U, Hagen S, Wist E, Moum T, and Kvikstad A. Psychological distress in cancer patients with advanced disease. Radiother Oncol 93;27(3):193-7. [Rec#: 3687]
INCLUDE: considered for evidence table

Kaldjian LC, Jekel JF, and Friedland G. End-of-life decisions in HIV-positive patients: the role of spiritual beliefs. AIDS 98;12(1):103-7. [Rec#: 12530]
INCLUDE: considered for evidence table

Kalra L, Evans A, Perez I, Knapp M, Donaldson N, and Swift CG. Alternative strategies for stroke care: a prospective randomised controlled trial. Lancet 2000;356(9233):894-9. Comment in: Lancet. 2000 Sep 9;356(9233):869-70. PMID: 11036886. Comment in: Lancet. 2001 Jan 13;357(9250):146-7. PMID: 11197428. Comment in: Lancet. 2001 Jan 13;357(9250):146; author reply 147. PMID: 11197429. [Rec#: 3712]
INCLUDE: considered for evidence table

Kalso E, Heiskanen T, Rantio M, Rosenberg PH, and Vainio A. Epidural and subcutaneous morphine in the management of cancer pain: a double-blind cross-over study. Pain 96;67(2-3):443-9. [Rec#: 3713]
INCLUDE: considered for evidence table

Kane RL, Homyak P, Bershadsky B, and Lum YS. Consumer responses to the Wisconsin Partnership Program for Elderly Persons: a variation on the PACE Model. J Gerontol A Biol Sci Med Sci 2002;57(4):M250-8. [Rec#: 41719]
INCLUDE: considered for evidence table

Kane RL, Keckhafer G, Flood S, Bershadsky B, and Siadaty MS. The effect of Evercare on hospital use. J Am Geriatr Soc 2003;51(10):1427-34. [Rec#: 41721]
INCLUDE: considered for evidence table

Karetzky M, Zubair M, and Parikh J. Cardiopulmonary resuscitation in intensive care unit and non-intensive care unit patients. Immediate and long-term survival. Arch Intern Med 95;155(12):1277-80. Comment in: Arch Intern Med. 1996 Jan 22;156(2):215-6. PMID: 8546559. [Rec#: 41729]
INCLUDE: considered for evidence table

Karlin NJ and Retzlaff PD. Psychopathology in caregivers of the chronically ill: personality and clinical syndromes. Hosp J 95;10(3):55-61. [Rec#: 12550]
INCLUDE: considered for evidence table

Keay TJ, Taler GA, Fredman L, and Levenson SA. Assessing medical care of dying residents in nursing homes. Am J Med Qual 97;12(3):151-6. [Rec#: 3799]
INCLUDE: considered for evidence table

Keefe FJ, Ahles TA, Porter LS, Sutton LM, McBride CM, Pope MS, McKinstry ET, Furstenberg CP, Dalton J, and Baucom DH. The self-efficacy of family caregivers for helping cancer patients manage pain at end-of-life. Pain 2003;103(1-2):157-62. [Rec#: 3801]
INCLUDE: considered for evidence table

Keegan O, McGee H, Hogan M, Kunin H, O'Brien S, and O'Siorain L. Relatives' views of health care in the last year of life. Int J Palliat Nurs 2001;7(9):449-56. [Rec#: 12596]
INCLUDE: considered for evidence table

Keenan SP, Busche KD, Chen L. M, Esmail R, Inman KJ, and Sibbald WJ. Withdrawal and withholding of life support in the intensive care unit: a comparison of teaching and community hospitals. The Southwestern Ontario Critical Care Research Network. Critical Care Medicine 98;26(2):245-51. [Rec#: 50513]
INCLUDE: considered for evidence table

Keenan, SP, Busche, KD, Chen, LM, McCarthy, L, Inman, KJ, and Sibbald, WJ. A retrospective review of a large cohort of patients undergoing the process of withholding or withdrawal of life support. Critical Care Medicine 97;25(8):1324-31. [Rec#: 50514]
INCLUDE: considered for evidence table

Appendix M. Included Studies

Keene J, Hope T, Fairburn CG, and Jacoby R. Death and dementia. Int J Geriatr Psychiatry 2001;16(10):969-74. Comment in: Int J Geriatr Psychiatry. 2002 Nov;17(11):1078. PMID: 12404661. [Rec#: 41748]
INCLUDE: considered for evidence table

Kehayia E, Korner-Bitensky N, Singer F, Becker R, Lamarche M, Georges P, and Retik S. Differences in pain medication use in stroke patients with aphasia and without aphasia. Stroke 97;28(10):1867-70. [Rec#: 3803]
INCLUDE: considered for evidence table

Kellogg FR and Ramos A. Code status decision-making in a nursing home population: processes and outcomes. J Am Geriatr Soc 95;43(2):113-21. [Rec#: 3815]
INCLUDE: considered for evidence table

Kelly B, Edwards P, Synott R, Neil C, Baillie R, and Battistutta D. Predictors of bereavement outcome for family carers of cancer patients. Psychooncology 99;8(3):237-49. [Rec#: 3819]
INCLUDE: considered for evidence table

Kennard MJ, Speroff T, Puopolo AL, Follen MA, Mallatratt L, Phillips R, Desbiens N, Califf RM, and Connors AF Jr. Participation of nurses in decision making for seriously ill adults. Clin Nurs Res 96;5(2):199-219. [Rec#: 3840]
INCLUDE: considered for evidence table

Kernerman P, Cook DJ, and Griffith LE. Documenting life-support preferences in hospitalized patients. J Crit Care 97;12(4):155-60. [Rec#: 41760]
INCLUDE: considered for evidence table

Kerridge IH, Pearson SA, Rolfe IE, and Lowe M. Decision making in CPR: attitudes of hospital patients and healthcare professionals. Med J Aust 98;169(3):128-31. Comment in: Med J Aust. 1998 Aug 3;169(3):124-5. PMID: 9734504. Comment in: Med J Aust. 1999 Jan 4;170(1):45-6. PMID: 10026675. Comment in: Med J Aust. 1999 Jan 4;170(1):45; author reply 45-6. PMID: 10026674. Comment in: Med J Aust. 1999 May 3;170. [Rec#: 3855]
INCLUDE: considered for evidence table

Kiely DK, Mitchell SL, Marlow A, Murphy KM, and Morris JN. Racial and state differences in the designation of advance directives in nursing home residents. J Am Geriatr Soc 2001;49(10):1346-52. [Rec#: 12673]
INCLUDE: considered for evidence table

Kimball LR and McCormick WC. The pharmacologic management of pain and discomfort in persons with AIDS near the end of life: use of opioid analgesia in the hospice setting. J Pain Symptom Manage 96;11(2):88-94. [Rec#: 3892]
INCLUDE: considered for evidence table

Kirschling JM, Stewart BJ, and Archbold PG. Family caregivers of post-hospitalized older persons and persons receiving hospice: similarities and differences. Home Health Care Serv Q 94;14(4):117-40. [Rec#: 3923]
INCLUDE: considered for evidence table

Kish Wallace S, Martin CG, Shaw AD, and Price KJ. Influence of an advance directive on the initiation of life support technology in critically ill cancer patients. Crit Care Med 2001;29(12):2294-8. Comment in: Crit Care Med. 2001 Dec;29(12):2391-2. PMID: 11801850. [Rec#: 41781]
INCLUDE: considered for evidence table

Kissane DW, McKenzie M, McKenzie DP, Forbes A, O'Neill I, and Bloch S. Psychosocial morbidity associated with patterns of family functioning in palliative care: baseline data from the Family Focused Grief Therapy controlled trial. Palliat Med 2003;17(6):527-37. [Rec#: 3934]
INCLUDE: considered for evidence table

Knol HR, Haken L, and Kempen GI. Disablement process and the utilization of home care among non-institutionalized elderly people: contrasting results between cross-sectional and panel data. Disabil Rehabil 2003;25(15):845-55. [Rec#: 41806]
INCLUDE: considered for evidence table

Koeberle D, Bacchus L, Thuerlimann B, and Senn HJ. Pamidronate treatment in patients with malignant osteolytic bone disease and pain: a prospective randomized double-blind trial. Support Care Cancer 99;7(1):21-7. Comment in: Support Care Cancer. 1999 Jan;7(1):1-3. PMID: 9926966. [Rec#: 3980]
INCLUDE: considered for evidence table

Koedoot CG, Oort FJ, de Haan RJ, Bakker PJ, de Graeff A, and de Haes JC. The content and amount of information given by medical oncologists when telling patients with advanced cancer what their treatment options are. Palliative chemotherapy and watchful-waiting. Eur J Cancer 2004;40(2):225-35. [Rec#: 3984]
INCLUDE: considered for evidence table

Koenig HG. Depression in hospitalized older patients with congestive heart failure. Gen Hosp Psychiatry 98;20(1):29-43. [Rec#: 41814]
INCLUDE: considered for evidence table

Koffman J, Higginson IJ, and Donaldson N. Symptom severity in advanced cancer, assessed in two ethnic groups by interviews with bereaved family members and friends. J R Soc Med 2003;96(1):10-6. [Rec#: 12762]
INCLUDE: considered for evidence table

Kolarik RC, Arnold RM, Fischer GS, and Hanusa BH. Advance care planning. J Gen Intern Med 2002;17(8):618-24. [Rec#: 3998]
INCLUDE: considered for evidence table

Kongsgaard UE and Meidell NK. Mechanical ventilation in critically ill cancer patients: outcome and utilisation of resources. Support Care Cancer 99;7(2):95-9. [Rec#: 4010]
INCLUDE: considered for evidence table

Kornblith AB, Herndon JE, Zuckerman E, Viscoli CM, Horwitz RI, Cooper MR, Harris L, Tkaczuk KH, Perry MC, Budman D, Norton LC, Holland J, Breast Cancer and Leukemia Group. Social support as a buffer to the psychological impact of stressful life events in women with breast cancer. Cancer 2001;91(2):443-54. [Rec#: 50526]
INCLUDE: considered for evidence table

Kramer BJ and Kipnis S. Eldercare and work-role conflict: toward an understanding of gender differences in caregiver burden. Gerontologist 95;35(3):340-8. [Rec#: 41841]
INCLUDE: considered for evidence table

Kristjanson LJ, Leis A, Koop PM, Carriere KC, and Mueller B. Family members' care expectations, care perceptions, and satisfaction with advanced cancer care: results of a multi-site pilot study. J Palliat Care 97;13(4):5-13. [Rec#: 4063]
INCLUDE: considered for evidence table

Kristjanson LJ, Sloan JA, Dudgeon D, and Adaskin E. Family members' perceptions of palliative cancer care: predictors of family functioning and family members' health. J Palliat Care 96;12(4):10-20. [Rec#: 4066]
INCLUDE: considered for evidence table

Kristjanson, L. J. Validity and reliability testing of the FAMCARE Scale: measuring family satisfaction with advanced cancer care. Social Science & Medicine 93;36(5):693-701. [Rec#: 50531]
INCLUDE: considered for evidence table

Kristjansson B, Breithaupt K, and McDowell I. Development and validation of an indicator of support for community-residing older Canadians. Int Psychogeriatr 2001;13 Supp 1:125-35. [Rec#: 41855]
INCLUDE: considered for evidence table

Krout RE. The effects of single-session music therapy interventions on the observed and self-reported levels of pain control, physical comfort, and relaxation of hospice patients. Am J Hosp Palliat Care 2001;18(6):383-90. [Rec#: 4074]
INCLUDE: considered for evidence table

Krumholz HM, Phillips RS, Hamel MB, Teno JM, Bellamy P, Broste SK, Califf RM, Vidaillet H, Davis RB, Muhlbaier LH, Connors AF Jr, Lynn J, and Goldman L. Resuscitation preferences among patients with severe congestive heart failure: results from the SUPPORT project. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. Circulation 98;98(7):648-55. Comment in: Circulation. 1998 Aug 18;98(7):619-22. PMID: 9715851. [Rec#: 4078]
INCLUDE: considered for evidence table