

**A cohort-study of patients suspected for MERS-CoV in a referral hospital in Saudi Arabia**

Jaffar A. Al-Tawfiq <sup>a,b</sup>, Sarah H. Alfaraj<sup>c</sup>, Talal A. Altuwaijri <sup>d</sup>, Ziad A. Memish <sup>e, f,g</sup>

<sup>a</sup> Johns Hopkins Aramco Healthcare, Dhahran, Kingdom of Saudi Arabia

<sup>b</sup> Indiana University School of Medicine, Indianapolis, IN, USA

<sup>c</sup> Corona Center, Infectious Diseases Division, Department of Pediatric , Prince Mohamed Bin Abdulaziz Hospital, Ministry of Health, Riyadh, Saudi Arabia

<sup>d</sup> Department of Surgery, Prince Mohamed Bin Abdulaziz Hospital, Ministry of Health, Riyadh, Saudi Arabia

<sup>e</sup> College of Medicine, Alfaisal University, Riyadh, Saudi Arabia

<sup>f</sup>. Infectious Diseases Division, Department of Medicine, Prince Mohamed Bin Abdulaziz Hospital, Ministry of Health, Riyadh, Saudi Arabia

<sup>g</sup> Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, USA

**All authors have no conflict of interest to report**

---

This is the author's manuscript of the article published in final edited form as:

Al-Tawfiq, J. A., Alfaraj, S. H., Altuwaijri, T. A., & Memish, Z. A. (2017). A cohort-study of patients suspected for MERS-CoV in a referral hospital in Saudi Arabia. *Journal of Infection*.  
<https://doi.org/10.1016/j.jinf.2017.06.002>

Ko and colleagues recently published predictive factors for pneumonia in patients infected with Middle East respiratory syndrome-coronavirus (MERS-CoV) [1]. The differentiation between MERS-CoV and non-MERS-CoV was evaluated in few studies with no significant findings [2–4]. In this study, we review a large cohort of patients who were admitted to one (Prince Mohammed bin Abdulaziz Hospital) of the 4 MERS-CoV reference centers under the umbrella of the Ministry of Health (MoH) to rule out MERS-CoV with the aim to compare MERS-CoV positive patients to non-MERS-CoV patients. Prince Mohammed bin Abdulaziz Hospital (PMAH) is referral center for all MERS-CoV patients diagnosed in the central region based in Riyadh, Saudi Arabia. The study included all patients from April 1<sup>st</sup> 2014 to December 2016. MERS-CoV testing was done using nasopharyngeal swabs and tested using real time reverse transcriptase PCR as described previously [5]. In this cohort, we compared MERS-CoV patients to non-MERS-CoV patients. The comparison used percentage for categorical data and mean  $\pm$  standard deviation (SD) for continuous data. A p value of  $< 0.05$  was considered significant.

During the study period, there were a total of 2151 patients who were admitted for evaluation for MERS-CoV. Of those patients, 1856 (86.3%) were non-MERS-CoV, and 295 (13.7%) were MERS-CoV patients. Male gender was 61% among MERS-CoV patients and 53.7% among non-MERS-CoV patients ( $P = 0.019$ ). There was no difference in the mean age of MERS-CoV patients ( $47.3 \pm 17.8$  years) compared with  $554.6 \pm 22.1$  years in the non-MERS-CoV patients. Overall, the age group 21-40 and 41-60 years were more common among MERS-CoV patients (36.9% vs. 24.4% and 32.9% vs. 22.5%, respectively). Cough and shortness of breath were less common in the MERS-CoV than non-MERS-CoV patients (41.4% vs. 50.3% and 54.2% vs. 66%). Healthcare workers represented 14.9% and 2.6% ( $P = 0.0001$ ) in the MERS-CoV than non-MERS-CoV patients, respectively. The mean WBC count was not significantly lower in

MERS-CoV vs. non-MERS-CoV patients (mean 7.47 vs 11.5,  $P = 1$ ). The case fatality rate was significantly lower in the non-MERS-CoV (9.2%) compared with MERS-CoV (20%) ( $P = 0.0001$ ).

This is the largest cohort study of MERS-CoV and non-MERS-CoV patients. Previously, there were three case-control studies of MERS-CoV vs. non-MERS-CoV patients. The first study included 17 MERS-CoV patients and 82 non-MERS-CoV patients with a high mortality rate of 76% and 15% among MERS-CoV and non-MERS-CoV patients respectively. The second study included 80 MERS-CoV patients and was matched to 159 non-MERS-CoV patients with a recorded mortality rate of 10% vs. 4.4%. A third study from Saudi Arabia included 48 patients with MERS-CoV infection and 111 MERS-CoV negative patients. The study included patients from October 2012 to April 2014 and the case fatality rates were 35% and 22% respectively. The current study included a larger number of patients spanning a longer duration and the case fatality rate was only 20%. This case fatality rate was different in various hospitals and ranged from 28.6% to 63.6% [6]. The current study had a case fatality rate that is close to the South Korea rate and lower than the overall all rate (38.7%) and the Kingdom-wide rate of 36.5% [7]. The variability of the case-fatality rates may be related to host factors, co-morbidities, care provided and yet unidentified factors [6].

Most of the cases of MERS occurred in the healthcare settings [2,8]. The proportion of healthcare workers (HCWs) in these outbreaks is variable. In a study of 70 cases, at least 50% of the cases were acquired in the hospital setting [9]. In the 2014 Jeddah outbreak, there were 128 laboratory-confirmed cases in 14 hospitals between February 17 and April 26, 2014 [10]. Of these cases, 33% were primary cases and >60% (including 39 HCWs) were acquired within healthcare facilities. In the current study, HCWs represented 14.9% and 2.6% ( $P = 0.0001$ ) in

the MERS-CoV and non-MERS-CoV patients, respectively. However, the hospital is a referral hospital for all MERS cases in the Riyadh region and thus the finding does not represent the transmission within the same hospital. In conclusion, the case fatality rate was higher in the MERS than non-MERS and healthcare workers were more common among MERS patients.

## References:

- [1] Ko J-H, Park GE, Lee JY, Lee JY, Cho SY, Ha YE, et al. Predictive factors for pneumonia development and progression to respiratory failure in MERS-CoV infected patients. *J Infect* 2016;73:468–75. doi:10.1016/j.jinf.2016.08.005.
- [2] Al-Tawfiq JA, Hinedi K, Ghandour J, Khairalla H, Musleh S, Ujayli A, et al. Middle East Respiratory Syndrome-Coronavirus (MERS-CoV): a case-control study of hospitalized patients. *Clin Infect Dis* 2014;59:160–5. doi:10.1093/cid/ciu226.
- [3] Mohd HA, Memish ZA, Alfaraj SH, McClish D, Altuwaijri T, Alanazi MS, et al. Predictors of MERS-CoV infection: A large case control study of patients presenting with ILI at a MERS-CoV referral hospital in Saudi Arabia. *Travel Med Infect Dis* 2016;14:464–70. doi:10.1016/j.tmaid.2016.09.008.
- [4] Garbati MA, Fagbo SF, Fang VJ, Skakni L, Joseph M, Wani TA, et al. A Comparative Study of Clinical Presentation and Risk Factors for Adverse Outcome in Patients Hospitalised with Acute Respiratory Disease Due to MERS Coronavirus or Other Causes. *PLoS One* 2016;11:e0165978. doi:10.1371/journal.pone.0165978.
- [5] Assiri A, Al-Tawfiq JA, Al-Rabeeh AA, Al-Rabiah FA, Al-Hajjar S, Al-Barrak A, et al.

- Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: A descriptive study. *Lancet Infect Dis* 2013;13:752–61. doi:10.1016/S1473-3099(13)70204-4.
- [6] Nam H-S, Park JW, Ki M, Yeon M-Y, Kim J, Kim SW. High fatality rates and associated factors in two hospital outbreaks of MERS in Daejeon, the Republic of Korea. *Int J Infect Dis* 2017. doi:10.1016/j.ijid.2017.02.008.
- [7] Kim KH, Tandi TE, Choi JW, Moon JM, Kim MS. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak in South Korea, 2015: epidemiology, characteristics and public health implications. *J Hosp Infect* 2017;95:207–13. doi:10.1016/j.jhin.2016.10.008.
- [8] Al Hosani FI, Pringle K, Al Mulla M, Kim L, Pham H, Alami NN, et al. Response to Emergence of Middle East Respiratory Syndrome Coronavirus, Abu Dhabi, United Arab Emirates, 2013–2014. *Emerg Infect Dis* 2016;22:1162–8. doi:10.3201/eid2207.160040.
- [9] Saad M, Omrani AS, Baig K, Bahloul A, Elzein F, Matin MA, et al. Clinical aspects and outcomes of 70 patients with Middle East respiratory syndrome coronavirus infection: a single-center experience in Saudi Arabia. *Int J Infect Dis* 2014;29:301–6. doi:10.1016/j.ijid.2014.09.003.
- [10] Drosten C, Muth D, Corman VM, Hussain R, Al Masri M, HajOmar W, et al. An observational, laboratory-based study of outbreaks of middle East respiratory syndrome coronavirus in Jeddah and Riyadh, kingdom of Saudi Arabia, 2014. *Clin Infect Dis* 2015;60:369–77. doi:10.1093/cid/ciu812.

**Table 1: Comparison of characteristics of non-MERS-CoV and MERS-CoV patients**

	<b>Non MERS-CoV (Number)</b>	<b>Non-MERS-CoV (%)</b>	<b>MERS-CoV (Number)</b>	<b>MERS-CoV (%)</b>	<b>P value</b>
<b>Number</b>	1856		295		
<b>Male</b>	997	53.7	180	61.0	0.0199
<b>0-20</b>	116	6.3	8	2.7	0.014
<b>21-40</b>	453	24.4	109	36.9	0.0001
<b>41-60</b>	417	22.5	97	32.9	0.0002
<b>61-80</b>	683	36.8	75	25.4	0.0001
<b>81-100</b>	187	10.1	6	2.0	0.0001
<b>Abnormal</b>					
<b>CXR</b>	1170	63.0	182	61.7	0.6505
<b>Fever</b>	1718	92.6	282	95.6	0.065
<b>Symptoms</b>	1814	97.7	252	85.4	0.0001
<b>GI symptoms</b>	228	12.3	32	10.8	0.5638
<b>SOB</b>	934	50.3	122	41.4	0.0047
<b>Cough</b>	1225	66.0	160	54.2	0.0001
<b>Healthcare</b>					
<b>workers</b>	48	2.6	44	14.9	0.0001
<b>Mechanical</b>					
<b>ventilation</b>	168	9.1	69	23.4	0.0001
<b>Death</b>	171	9.2	59	20.0	0.0001

**Table 2: Comparison of the means between non-MERS-CoV and MERS-CoV patients**

	<b>Non-MERS-CoV</b>	<b>SD</b>	<b>MERS-CoV</b>	<b>SD</b>	<b>P value</b>
<b>Mean age</b>	54.6	22.1	47.3	17.8	1.0
<b>WBC</b>	11.5	6.53	7.47	4.31	1.0
<b>Hgb</b>	125.8	27.1	129.1	25.1	0.037
<b>Platelet</b>	280	131.4	227	107.7	< 0.0001
<b>PMN Absolute</b>	14.72	128.0	9.46	65.0	0.29